

British Columbia (BC) COVID-19 Situation Report

Week 27: July 03- July 09, 2022

Data for week 27 (July 03 - July 09, 2022) may differ from the data published in the BCCDC weekly report. Data was extracted on July 18, 2022 for this situation report compared to July 20, 2022 for the latest weekly report.

Table of Contents	Report Summary
Epidemic curve and regional incidence 3	Due to changes in testing strategies in BC, current case counts are an underestimate of the true number of COVID-19 cases in BC. This underestimation has increased compared to the period prior to the emergence of the Omicron variant in BC. The provincial incidence by episode date was 19 per 100K (974 cases) in week 27, which increased from 15 per 100K in week 26.
Test rates and % positive 4	Incidence by Health Authority from week 26 to week 27: <ul style="list-style-type: none"> Fraser Health incidence increased from 17 to 20 per 100K Interior Health incidence remained stable at 12 and 13 per 100K Vancouver Island Health incidence increased from 15 to 22 per 100K Northern Health incidence increased from 12 to 20 per 100K Vancouver Coastal Health incidence increased from 13 to 17 per 100K
Age profile, testing and cases 5	Testing of MSP-funded specimens increased from ~6,400 in week 26 to ~6,750 in week 27, and the percent positivity of MSP-funded specimens increased from 14.0% in week 26 to 16.3% in week 27.
Severe outcomes 7	The per capita testing rates for MSP-funded specimens between week 26 and week 27 increased in all age groups except in the 0-4 and 5-9 year-olds, where testing rates decreased from 172 per 100K in week 26 to 155 per 100K in week 27 and from 69 per 100K in week 26 to 49 per 100K in week 27, respectively. Percent positivity between week 26 and week 27 increased or remained stable in all age groups except in the 10-14 year-olds where percent positivity decreased from 10.6% in week 26 to 8.6% in week 27.
Age profile, severe outcomes 9	Age-specific incidence rates between week 26 and week 27 increased or remained stable in all age groups. Incidence rates increased the most in the 80+ year-olds from 84 per 100K in week 26 to 104 per 100K in week 27.
Care facility outbreaks 11	The number of people in hospital with a positive COVID-19 test remained stable at 222 in week 26 and 225 in week 27. In week 27, 60+ year-olds had the highest number of people in hospital with a positive COVID-19 test, with 85 hospitalizations in 60-79 year-olds and 78 hospitalizations in 80+ year-olds.
Wastewater surveillance 12	The weekly number of deaths from any cause among people testing positive for COVID-19 decreased from 35 in week 26 to 28 in week 27. Similar to previous weeks, 80+ year-olds had the highest number of deaths from any cause among people testing positive for COVID-19 (17 deaths) in week 27. From week 14 to week 22 where the underlying cause of death (UCD) has been reported for at least 95% of the post-transition deaths, an average of 44% of these deaths were reported to have COVID-19 as their UCD.
Additional resources 14	In week 27, based on earliest symptom onset date (if unavailable, then outbreak declared date is used), 4 new care facility outbreaks were declared in long-term care.
Appendix 14	

BELOW ARE IMPORTANT NOTES relevant to the interpretation of cases, hospitalizations, and deaths:

- Due to changes in testing strategies in BC in 2022 focusing on targeted higher risk populations, current case counts are an underestimate of the true number of COVID-19 cases in BC. This underestimation has increased compared to the period prior to the emergence of the Omicron variant in BC.
- Hospital data include admissions for people who test positive for COVID-19 through hospital screening practices, regardless of the reason for admission. Therefore, reported hospitalizations overestimate the true number of people who are hospitalized specifically due to COVID-19 infection.
- Pre-transition (case line list) deaths include COVID-19 related deaths reported by Health Authorities up to April 1, 2022. As of April 2, 2022, post-transition (automated linkage) deaths include people who died from any cause recorded in Vital Statistics within 30 days of their first positive COVID-19 lab result date. Deaths reported after the system transition use a broader definition and will overestimate the true number of deaths due to COVID-19 since death registration is recorded before the underlying cause of death is determined. Due to the change in data source for death data, the number of pre-transition deaths should not be compared to the number of post-transition deaths.

BELOW ARE IMPORTANT NOTES relevant to the interpretation of data displayed in this bulletin:

- Cases include lab confirmed, lab probable, and epi-linked cases. Case definition can be found at [http://www.bccdc.ca/health-professionals/clinical-resources/case-definitions/covid-19-\(novel-coronavirus\)](http://www.bccdc.ca/health-professionals/clinical-resources/case-definitions/covid-19-(novel-coronavirus)). Cases include those reported in Health Authority case line lists and positive laboratory results in the Provincial Laboratory Information Solution (PLIS) up to April 1, 2022. As of April 2, 2022, only positive laboratory results in the PLIS are included and cases who are residents from outside of BC are not included.
 - Episode date is defined by date of illness onset when available. When illness onset date is unavailable, earliest laboratory date is used (collection or result date); if also unavailable, then public health case report date is used. As of April 2, 2022, episode date reflects earliest laboratory date (collection or result date) only. Analyses based on episode date may better represent the timing of epidemic evolution. Episode-based tallies for recent weeks are expected to increase as case data are more complete.
 - Surveillance date is defined by lab result date, if unavailable, then public health case report date is used. As of April 2, 2022, surveillance date reflects lab result date only. The weekly tally by surveillance date includes cases with illness onset date in preceding weeks.
 - Hospitalizations include those reported by Health Authorities up to April 1, 2022. As of April 2, 2022, hospitalizations are defined as individuals who test positive for COVID-19 and are hospitalized as recorded in the PHSA Provincial COVID-19 Monitoring Solution (PCMS). Hospitalizations for individuals 0-19 years-old are reported by linked hospitalization episodes from the PCMS since the beginning of the pandemic. Episode date for hospitalization is defined by admission date, if unavailable, surveillance date is used.
 - Critical care admissions (HAU, ICU, and critical care surge beds) include individuals who test positive for COVID-19 and are in critical care admission as recorded in the PCMS. Episode date for critical care admission is defined by critical care admission date, if unavailable, surveillance date is used. Previously only ICU admissions were presented in this report. Critical care admissions comprises a broader category than ICU admissions and therefore, the number of critical care admissions should not be compared to number of ICU admissions from previous weeks.
 - Deaths include COVID-19 related deaths reported by Health Authorities up to April 1, 2022. As of April 2, 2022, deaths are any COVID-19 lab positive cases who died from any cause recorded in Vital Statistics within 30 days of their first positive lab result date. Episode date for death is defined by death date, if unavailable, surveillance date is used.
 - As of April 2, 2022, data on Health Authority outbreaks are compiled from outbreak files provided by the Health Authorities.
 - Laboratory PLOVER data include Medical Service Plan (MSP) funded (e.g. clinical diagnostic tests) and non-MSP funded (e.g. screening tests) specimens.
 - Per capita rates/incidences for year 2020 are based on Population Estimates 2020 (n= 5,147,772 for BC overall), for year 2021 are based on PEOPLE 2021 estimates (n= 5,194,137 for BC overall), and for year 2022 is based on PEOPLE 2021 estimates (n= 5,263,772 for BC overall).
 - Data sources include Health Authority case line lists, PHSA Provincial COVID-19 Monitoring Solution (PCMS), Vital Statistics, laboratory PLOVER data, and aggregate outbreak files from Health Authorities.
 - Integrated case data (including surveillance variables created using Health Authority case line lists, PCMS, and Vital Statistics) were extracted on July 18, 2022, laboratory PLOVER data on July 14, 2022, and Health Authority outbreak files on July 13, 2022.
-

A. COVID-19 case counts and epidemic curves

Due to changes in testing strategies in BC in 2022 focusing on targeted higher risk populations, current case counts are an underestimate of the true number of COVID-19 cases in BC. This underestimation has increased compared to the period prior to the emergence of the Omicron variant in BC. Up to week 27, there have been 376,391 cases for a cumulative incidence of 7,151 per 100K (Table 1, Figure 1). The provincial incidence by episode date was 19 per 100K (974 cases) in week 27, which increased from 15 per 100K in week 26. Incidence by episode date may increase as data become more complete in recent weeks.

As shown in Figure 2, incidence rates from week 26 to week 27 increased or remained stable in all HAs. Incidence rates increased the most in the Northern Health (NH) from 12 per 100K in week 26 to 20 per 100K in week 27. In week 27, the highest incidence rate was in Island Health (VIHA) at 22 per 100K.

Table 1. Episode-based case tallies by Health Authority, BC, Jan 15, 2020 (week 3) – Jul 09, 2022 (week 27) (N= 376,391)

Case tallies by episode date	Health Authority of Residence					Outside Canada	Total
	FH	IH	VIHA	NH	VCH		
Week 27, case counts	395	106	193	62	218	0	974
Cumulative case counts	165,797	66,935	36,819	30,546	75,903	391	376,391
Week 27, cases per 100K population	20	13	22	20	17	NA	19
Cumulative cases per 100K population	8,343	8,080	4,183	9,980	6,015	NA	7,151

Figure 1. Episode-based epidemic curve (bars), surveillance date (line) and Health Authority (HA), BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27) (N= 368,544)





Figure 2. Weekly episode-based incidence rates by HA and health service delivery area (HSDA), BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27) (N= 368,544)



B. Test rates and percent positive

[COVID-19 testing guidelines](#) recommend testing for people who have COVID-19 symptoms, and are at risk of more severe disease or live/work in high-risk settings. As shown by the darker-colored bars and dotted line in [Figure 3](#), the number of MSP-funded specimens increased from ~6,400 in week 26 to ~6,750 in week 27, and the percent positivity of MSP-funded specimens increased from 14.0% in week 26 to 16.3% in week 27.

As shown by the dotted lines in [Figure 4](#), the per capita testing rates for MSP-funded specimens (dotted lines in Panel A) increased in all HAs except IHA, where it decreased from 88 per 100K in week 26 to 82 per 100K in week 27. In week 27, FH had the highest testing rate at 167 per 100K. The percent positivity (dotted lines in Panel B) for MSP-funded specimens increased or remained stable in all HAs. In week 27, percent positivity ranged from 13.5% in FH to 24.0% in VIHA.

Figure 3. Number of specimens tested and percent SARS-CoV-2 positive, by collection week, BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27)





Figure 4. Testing rates and percent SARS-CoV-2 positive by Health Authority and collection week, BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27)



Data source: Laboratory PLOVER data

C. Age profile – Testing and cases

Testing rates and percent positivity by age group


As shown by the bars in [Figure 5](#), the per capita testing rates for MSP-funded specimens between week 26 and week 27 increased in all age groups except in the 0-4 and 5-9 year-olds, where testing rates decreased from 172 per 100K in week 26 to 155 per 100K in week 27 and from 69 per 100K in week 26 to 49 per 100K in week 27, respectively.

As shown by the black dots in [Figure 5](#), percent positivity between week 26 and week 27 increased or remained stable in all age groups except in the 10-14 year-olds where percent positivity decreased from 10.6% in week 26 to 8.6% in week 27. Percent positivity increased the most in 0-4 and 60-79 year-olds, where it increased from 12.0% in week 26 to 16.0% in week 27 and from 12.2% in week 26 to 15.7% in week 27, respectively. In week 27, percent positivity ranged from 8.4% in 5-9 year-olds to 20.5% in 80+ year-olds.

Case distribution and weekly incidence by age group


As shown in [Figure 6](#), age-specific incidence rates between week 26 and week 27 increased or remained stable in all age groups. Incidence rates increased the most in the 80+ year-olds from 84 per 100K in week 26 to 104 per 100K in week 27.

Figure 5. Average weekly SARS-CoV-2 MSP testing rates and MSP percent positive by known age group, BC Jun 04, 2022 (week 22) – Jul 09, 2022 (week 27)



Data source: Laboratory PLOVER data

Figure 6. Weekly age-specific COVID-19 incidence per 100K population by epidemiological week, BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27) (N= 368,450)



D. Severe outcome counts and epi-curve

Hospital data include admissions for people who test positive for COVID-19 through hospital screening practices, regardless of the reason for admission. Therefore, reported hospitalizations overestimate the true number of people who are hospitalized specifically due to COVID-19 infection. The number of people in hospital with a positive COVID-19 test remained stable at 222 in week 26 and 225 in week 27. In week 27, 60+ year-olds had the highest number of people in hospital with a positive COVID-19 test, with 85 hospitalizations in 60-79 year-olds and 78 hospitalizations in 80+ year-olds.

As of April 2, 2022, death data include people who test positive for COVID-19 and died from any cause (COVID-19 or non-COVID-19) within 30 days of their first positive lab result date. The weekly number of deaths from any cause among people testing positive for COVID-19 decreased from 35 in week 26 to 28 in week 27. Similar to previous weeks, 80+ year-olds had the highest number of deaths from any cause among people testing positive for COVID-19 (17 deaths) in week 27 ([Table 2, Figure 8](#)). Detailed information about outcomes by vaccination status can be accessed at [BCCDC COVID-19 Regional Surveillance Dashboard](#).


Cumulatively, there have been 32 confirmed cases of [Multi-system Inflammatory Syndrome in children and adolescents \(MIS-C\)](#) in BC since January 1, 2020. There have been no new confirmed cases of MIS-C since the last report. The median age of all cases is 9 years old (range from 4 months old to 16 years old).

**Table 2. COVID-19 severe outcomes by episode date, Health Authority of residence, BC
Jan 15, 2020 (week 3) – Jul 09, 2022 (week 27)**

Severe outcomes by episode date	Health Authority of residence					Residing outside of Canada	Total n/N ^a (%)
	FH	IH	VIHA	NH	VCH		
Week 27, hospitalizations	106	21	33	12	53	0	225
Cumulative hospitalizations	11,581	4,194	2,402	2,084	4,926	17	25,204/376,391 (7)
Week 27, critical care admissions ^b	25	4	3	2	10	0	44
Cumulative critical care admissions^b	2,442	977	409	789	1,088	4	5,709/376,391 (2)
Week 27, deaths	16	3	4	2	3	0	28
Cumulative deaths, pre-transition (case line list)^c	1,348	367	241	330	716	0	3,002/356,546 (1)
Cumulative deaths, post-transition (automated linkage)^c	280	177	156	30	186	0	829/19,845 (4)

- Cases with unknown outcome are included in the denominators (i.e. assumed not to have the specified severe outcome).
- Due to the change in data source for hospitalization data, ICU admissions are no longer available. Critical care admissions are now being provided, which comprises a broader category than ICU admissions (please see Important Notes on Page 2 for more information). Number of critical care admissions should not be compared to number of ICU admissions from previous weeks.
- Pre-transition (case line list) deaths include COVID-19 related deaths reported by Health Authorities up to April 1, 2022. As of April 2, 2022, post-transition (automated linkage) deaths are any COVID-19 lab positive cases who died from any cause recorded in Vital Statistics within 30 days of their first positive lab result date. Deaths reported after the system transition use a broader definition and will overestimate the true number of deaths due to COVID-19 since death registration is recorded before the underlying cause of death is determined. Due to the change in data source for death data, the number of pre-transition deaths should not be compared to the number of post-transition deaths.

Figure 8. Weekly COVID-19 hospital admissions (A) and deaths (B) by age groups, BC, Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27)^a



a. Among those with available age information only.

E. Age profile, severe outcomes

Table 3 displays the distribution of cases and severe outcomes. In week 27, median age of hospital admissions, critical care admissions, pre-transition deaths, and post-transition deaths with underlying cause of death (UCD) as COVID-19 was 66 years, 63 years, 82 years, and 86 years, respectively.

In the past four weeks (from week 24 to week 27), there has been a weekly average of 3 deaths in those <60 years of age, 3 deaths in 60-69 year-olds, 8 deaths in 70-79 year-olds and 20 deaths in the 80+ year-olds (data not shown). The number of deaths may increase over time as data becomes more complete.


Table 3: COVID-19 cases, hospitalizations, critical care admissions, and deaths by age group, BC, Jan 15, 2020 (week 3) – Jul 09, 2022 (week 27) (N= 376,360)^a

Age group (years)	Cases	Hospitalizations n (%)	Critical care admissions ^b n (%)	Pre-transition (case line list) deaths ^c n (%)	Post-transition (automated linkage) deaths ^c		
					UCD as COVID-19 ^d n (%)	UCD as non-COVID-19 ^d n (%)	UCD pending ^d n (%)
<10	30,656	537 (2)	67 (<1)	2 (<1)	1 (<1)	2 (<1)	0 (<1)
10-19	35,758	345 (1)	50 (<1)	0 (<1)	0 (<1)	1 (<1)	0 (<1)
20-29	73,211	1,343 (2)	201 (<1)	6 (<1)	0 (<1)	5 (<1)	2 (<1)
30-39	70,087	2,294 (3)	417 (1)	31 (<1)	2 (<1)	5 (<1)	0 (<1)
40-49	54,145	2,204 (4)	577 (1)	64 (<1)	1 (<1)	7 (<1)	0 (<1)
50-59	44,002	3,095 (7)	1,044 (2)	166 (<1)	3 (<1)	18 (1)	4 (<1)
60-69	30,410	4,210 (14)	1,407 (5)	353 (1)	23 (1)	40 (2)	8 (<1)
70-79	17,754	4,883 (28)	1,293 (7)	655 (4)	58 (2)	94 (3)	26 (1)
80-89	13,242	4,433 (33)	574 (4)	989 (10)	109 (3)	136 (4)	33 (1)
90+	7,095	1,860 (26)	79 (1)	736 (15)	113 (5)	119 (6)	19 (1)
Total	376,360	25,204	5,709	3,002	310	427	92
Median age	36	66	63	82	86	83	81

- Among those with available age information only.
- Due to the change in data source for hospitalization data, ICU admissions are no longer available. Critical care admissions are now being provided, which comprises a broader category than ICU admissions (please see Important Notes on Page 2 for more information). Number of critical care admissions should not be compared to number of ICU admissions from previous weeks.
- Pre-transition (case line list) deaths include COVID-19 related deaths reported by Health Authorities up to April 1, 2022. As of April 2, 2022, post-transition (automated linkage) deaths are any COVID-19 lab positive cases who died from any cause recorded in Vital Statistics within 30 days of their first positive lab result date. Deaths reported after the system transition use a broader definition and will overestimate the true number of deaths due to COVID-19 since death registration is recorded before the underlying cause of death is determined. Due to the change in data source for death data, the number of pre-transition deaths should not be compared to the number of post-transition deaths.
- Since underlying cause of death (UCD) takes approximately 8 weeks to be recorded, all-cause mortality is initially reported and then retrospective evaluations of underlying cause of death are provided here to better understand true COVID-19 mortality. UCD as COVID-19 are deaths that have been determined to be caused by COVID-19 in their Vital Stats record. UCD as non-COVID-19 are deaths that have been determined to be not attributable to COVID-19 in their Vital Stats record that are reported as deaths due to a lab positive COVID-19 test within 30 days of death. UCD pending are all post-transition deaths that do not yet have a recorded UCD.

Figure 9 displays the number of pre-transition deaths and post-transition deaths (i.e. people who test positive for COVID-19 and died from any cause within 30 days of their first positive lab result date) by underlying cause of death as recorded in Vital Statistics from week 8 to week 27 in 2022. From week 14 to week 22 where the UCD has been reported for at least 95% of the post-transition deaths, an average of 44% of these deaths had an UCD as COVID-19. Post-transition deaths with complete UCD are expected to increase over time.

Figure 9: Pre- and post-transition deaths by underlying cause of death, BC, Feb 20, 2022 (week 8) – Jul 09, 2022 (week 27)^{a,b}



Dotted line represents system transition date

- a. Pre-transition (case line list) deaths include COVID-19 related deaths reported by Health Authorities up to April 1, 2022. As of April 2, 2022, post-transition (automated linkage) deaths are any COVID-19 lab positive cases who died from any cause recorded in Vital Statistics within 30 days of their first positive lab result date. Deaths reported after the system transition use a broader definition and will overestimate the true number of deaths due to COVID-19 since death registration is recorded before the underlying cause of death is determined. Due to the change in data source for death data, the number of pre-transition deaths should not be compared to the number of post-transition deaths.
- b. Since underlying cause of death (UCD) takes approximately 8 weeks to be recorded, all-cause mortality is initially reported and then retrospective evaluations of underlying cause of death are provided here to better understand true COVID-19 mortality. UCD as COVID-19 are deaths that have been determined to be caused by COVID-19 in their Vital Stats record. UCD as non-COVID-19 are deaths that have been determined to be not attributable to COVID-19 in their Vital Stats record that are reported as deaths due to a lab positive COVID-19 test within 30 days of death. UCD pending are all post-transition deaths that do not yet have a recorded UCD.


F. Care facility outbreaks

As shown in [Table 4](#) and [Figure 10](#), 684 care facility (acute care and long-term care settings) outbreaks were reported in total in BC to the end of week 27. In week 27, based on earliest symptom onset date (if unavailable, then outbreak declared date is used), 4 new care facility outbreaks were declared in long-term care. In the past four weeks (from week 24 to week 27), there has been a weekly average of 3 care facility outbreaks.

Table 4. COVID-19 care facility^a outbreaks by earliest case onset^{b,c}, associated cases and deaths by episode date, BC Jan 15, 2020 (week 3) – Jul 09, 2022 (week 27) (N=684)^{d,e}

Care facility outbreaks and cases by episode date	Outbreaks	Cases			Deaths		
		Residents	Staff/other	Total	Residents	Staff/other	Total
Week 27, Care Facility Outbreaks	4	49	0	49	0	0	0
Cumulative, Care Facility Outbreaks	684	9,676	3,817	13,493	1,453	0	1,453

Figure 10. COVID-19 care facility^a, outbreaks by earliest case onset^{b,c}, facility type (A) and Health Authority (B), BC Sept 13, 2020 (week 38) – Jul 09, 2022 (week 27) (N=616)^{d,e}



- Case and death counts include PCR positive cases only for outbreaks in NHA and VIHA. Vancouver Coastal Health, Fraser Health Authority, and Interior Health Authority outbreaks may also include those diagnosed by rapid antigen tests or considered as suspect reinfection.
- Earliest dates of onset of outbreak cases are subject to change as investigations and data are updated. If unavailable, outbreak declared date is used.
- New outbreaks reported since the last report with an earliest case onset date (if unavailable, outbreak declared date is used) prior to the current reporting week will be included in the cumulative care facility outbreak total.
- Cases with unknown role are included in the case count for Staff/other.
- Data might be incomplete or vary from what was reported previously due to updates by Health Authorities.

G. Wastewater surveillance

The BCCDC and Metro Vancouver measure SARS-CoV-2 in wastewater at five wastewater treatment plants (treating wastewater from 50% of BC’s population). To account for changing wastewater volume due to rainfall or snowmelt, SARS-CoV-2 concentrations are normalized to wastewater flow. Normalized SARS-CoV-2 wastewater levels (measured as viral copies per day) are shown alongside incident COVID-19 cases in each wastewater catchment area in [Figure 10](#) and [Figure 11](#). The BCCDC’s test results are obtained from the liquid fraction of the wastewater sample. Other organizations, such as the National Microbiology Laboratory, test from the solid fraction of wastewater and therefore, their results are not directly comparable.

Key messages with results through to July 16, 2022:

- Generally, increases in SARS-CoV-2 viral loads in wastewater have slowed or reversed in the past one to two weeks. Further data are required before determining whether this trend is consistent with plateauing of COVID-19 incidence in Metro Vancouver, or short-term variability of these environmental measurements.
- Over the past one week, viral loads at Annacis Island (Fraser North and South) have decreased by 13%.
- Over the past two weeks, viral loads at Northwest Langley (Northwest Langley) have decreased by 39%.
- Over the past week, viral loads at Iona Island (Vancouver) have decreased by 36%.
- Over the past week, viral loads at Lulu Island (Richmond) have decreased by 13%.
- Over the past week, viral loads at Lions Gate (North Shore) have decreased by 58%.

Figure 11. Wastewater surveillance, FH





Figure 12. Wastewater surveillance, VCH



H. Additional resources

For maps and geographical distribution of cases and vaccinations, visit the BCCDC COVID-19 Regional Surveillance Dashboard here: <http://www.bccdc.ca/health-professionals/data-reports/covid-19-surveillance-dashboard>

Variant of concern (VOC) findings are available weekly here: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/data#variants>

For local, national, and global comparisons of BC to other jurisdictions on key epidemiological metrics, visit the BCCDC COVID-19 Epidemiology App here: https://bccdc.shinyapps.io/covid19_global_epi_app/

I. Appendix

[Vaccination phases](#) defined by vaccine eligibility of target populations in BC

Vaccination Phase 1 (December 2020 – February 2021)

Target populations include residents, staff and essential visitors to long-term care settings; individuals assessed and awaiting a long-term care placement; health care workers providing care for COVID-19 patients; and remote and isolated Indigenous communities.

Vaccination Phase 2 (February 2021 – April 2021)

Target populations include seniors, age ≥ 80 ; Indigenous peoples age ≥ 65 and Indigenous Elders; Indigenous communities; hospital staff, community general practitioners and medical specialists; vulnerable populations in select congregate settings; and staff in community home support and nursing services for seniors.

Vaccination Phase 3 (April 2021 – May 2021)

Target populations include people aged 60-79 years, Indigenous peoples aged 18-64 and people aged 16-74 who are clinically extremely vulnerable.

Vaccination Phase 4 (May 2021 – November 2021)

Target populations include everyone 12+ years. In September, third dose is available for people who are clinically extremely vulnerable.

Vaccination Phase 5 (November 2021 – February 2022)

Target populations include everyone 5+. Children aged 5-11 are eligible at the end of November. Everyone 18 and older will be invited to get a booster dose within 6-8 months of their second dose.

Vaccination Phase 6 (February 2022 – April 2022)

Target populations include everyone 5+. Everyone 12 and older will be invited to get a booster dose within 6-8 months of their second dose.

Vaccination Phase 7 (April 2022 – Present)

Target populations include everyone 5+. Everyone 12 and older will be invited to get a booster dose within 6-8 months of their second dose. People in long-term care, assisted living, seniors and Indigenous people can get a second booster 6 months after the date of the first booster.