

You need to:

- 1 Check if you may benefit from treatment
- 2 Self-isolate
- 3 Manage your symptoms
- 4 Report your result

Treatment for COVID-19

There are some treatments available for people who have tested positive for COVID-19 and who are at higher risk of serious illness and needing hospitalization.

If you test positive for COVID-19, check to see if you may benefit from treatment.

Visit gov.bc.ca/covidtreatments or call 1-888-268-4319

Self-isolate

Self-isolation means keeping away from others to help stop the spread of COVID-19. To learn more, visit: bccdc.ca/covid19selfisolation

People younger than 18 years and fully vaccinated people 18 years and older

5 days



- You have isolated for at least **5 days**
- Your fever is gone without medication
- Your symptoms have improved

People 18 years and older who are not fully vaccinated

10 days



- You have isolated for at least **10 days**
- Your fever is gone without medication
- Your symptoms have improved

Fully vaccinated means you received both doses of a 2-dose series (e.g. AstraZeneca, Pfizer-BioNTech, or Moderna vaccine) or it has been more than 14 days since you received a single dose of a 1-dose series (e.g. Janssen).

Continue to isolate for longer if you have a fever or are not feeling better. When you end isolation, you are unlikely to pass on COVID-19. However, it can take longer to recover from the illness. Most people feel better within two weeks. Some people with more severe symptoms can take 12 weeks or more to feel entirely better. If you are unsure or concerned, connect with your health care provider, call **8-1-1**, or go to an Urgent and Primary Care Centre to be assessed.

If you have returned from outside Canada and test positive for COVID-19, you need to follow the quarantine requirements set by the Federal government. For more information, visit travel.gc.ca/travel-covid.

Manage your symptoms

Most people can safely manage their symptoms at home by drinking plenty of fluids, resting, and using a humidifier (if you have access to one) or hot shower to ease a cough or sore throat. If you have a fever, you can use non-prescription medicine like acetaminophen (such as Tylenol) or ibuprofen (such as Advil) to help with some of the symptoms of COVID-19.

You can call 8-1-1 anytime to talk to a nurse at HealthLinkBC. This service is available in 130 languages. If your symptoms worsen, or if you do not improve after five or six days, call 8-1-1, your family doctor or an Urgent and Primary Care Centre (UPCC), so they can determine if you need to be assessed again.

Go to an emergency department or call 9-1-1 if you:

- Find it hard to breathe
- Have chest pain
- Can't drink anything
- Feel very sick
- Feel confused



Report your test result

It is important for public health to know your health history, where you may have been during your infectious period, such as where you work or live. The information you provide will help public health prioritize people who may benefit from public health follow up, and/or treatment for COVID-19.

Report your results at:

reportcovidresults.bccdc.ca



Vaccination after COVID-19

If you're not fully vaccinated or you have not received your booster, you should still get vaccinated after you have recovered and ended your self-isolation.

Vaccines boost your immunity and have shown to be highly effective in preventing serious illness and death, even after you have had a COVID-19 infection.

Learn more about vaccination:

bccdc.ca/covid19vaccine

