


Weekly update on Variants of Concern (VOC)

Dec 17, 2021

Of all positive samples sequenced* in epi week 48 (Nov 28 - Dec 04) in BC, ~ 100% were confirmed VOCs. Over time, the distribution of variants demonstrate the temporality and changing nature of circulating Variants of Concern (VOCs) as shown in Figure 1.

Data from epi week 48 may reflect partial data; estimates are expected to change as more specimens are received and sequenced.

Figure 1. Twenty most prevalent lineages in British Columbia, Jan 1 - Dec 04, 2021**




*Data from the PLOVER system at the BCCDC Public Health Lab.

**Dashed lines indicate the time of changes in whole genome sequencing (WGS) sampling strategy (epi week 22: all positive samples; epi week 36: transition from full sequencing to sequencing a subset of 10% of representative samples in addition to all targeted samples, while keeping a monthly census of all positive samples on the first week of the month; epi week 46: transition to WGS of all positive samples).


As shown in Figure 1, the main circulating variant is Delta, accounting for about 98.7% of positive specimens sequenced, with spatial variability of its different sub-lineages by HA (see Figure 2).

Figure 2. Ten most prevalent lineages in British Columbia by Health Authority, June 1 - Dec 04, 2021



Please note that the estimate of distribution of VOC lineage[#] (Figure 3) in BC for latest epi week 48 (Nov 28 - Dec 04) may change as more sequencing results are analyzed and given the lag in receipt of positive samples from frontline laboratories.

Figure 3. Estimated Sample prevalence[^] of VOCs by lineage by epi week of collection date, Jan 3 - Dec 04, 2021



[^] Sample prevalence is calculated as the rate of occurrence of a given VOC lineage per 100 positive lab samples. It was estimated from the proportion of presumptive VOC from screening and the proportion of confirmed VOC via sequencing (excluding outbreaks and targeted surveillance) until May 30th, 2021 when BC transitioned to WGS on all positive cases. From week 13 (March 28, 2021), VOC screening results with both E484K and N501Y mutations are assumed to be Gamma, given a very low prevalence of Beta in BC. As of week 22 (May 30, 2021), prevalence of VOC is estimated from sequencing results only.

Pink shaded box can reflect partial data due to a lag in receipt of positive samples from front line laboratories and turn around time of 7 to 11 days from sample collection to WGS analysis; estimates are expected to change as more specimens are received and sequenced.

Dash-dotted line indicates the time of transition to WGS of all positive samples on May 30, 2021 (epi week 22).

Dash line indicates the time of transition from full sequencing to sequencing only a subset of 10% of representative samples in addition to all targeted samples, while keeping a monthly census of all positive samples on the first week of the month, starting September 5, 2021 (epi week 36).


Dotted line indicates the time of transition to WGS of all positive samples on November 15, 2021 (epi week 46).

See appendix for the definitions of VOC lineages

Monitoring of Variants

As illustrated in Figure 4 below, BCCDC Public Health Laboratory is continuously monitoring for both VOCs and VOIs and it is tracking a regularly updated Variants Under Monitoring (VUMs). There are numerous VOIs, and they may not necessarily become VOCs. Once a VOI becomes a VOC, it will be added to our VOC reporting. Since September 2021 BC has adopted a new sampling strategy for sequencing to report on the provincial number of variants based on weekly point prevalence. To address the new Omicron (B.1.1.529 and BA.*) sequencing of all positives samples has resumed with retrospective specimens collected from November 15th 2021 onwards.

Figure 4. Overview of the screening and sequencing process applied to positive COVID-19 tests in BC, Dec 2021.




Please note the turnaround time sequencing which takes approximately 7-11 days, but it could also take longer if there are lab backlogs or if there are delays in receiving current positive samples from frontline laboratories.

Whole genome sequencing (WGS)

Whole genome sequencing (Illumina only) was performed on 103,605 specimens up to epi week 48 (Nov 28 - Dec 04) in BC, of which 83,573 came back as variants under closer observation. Figure 4 above illustrates BC's whole genome sequencing strategy of COVID samples.


The VOCs represent 97% of all the variants that were detected in the province (see [WGS frequency of lineages table](#) on BCCDC website). The Delta variant accounts for largest proportion of the VOCs and includes B.1.617.2, the parent lineage, and two sub-lineages AY.25 and AY.27 (Figure 5).

Figure 5. Cumulative Distribution of the three most prevalent Delta lineages



BCCDC Public Health Laboratory updates the lineage assignment tool (Pangolin), on an at least weekly basis, to reflect current lineage classification changes. Figure 6 below demonstrates how these updates affect changes in the Delta variant from one sub-lineage to another over time.

Figure 6. Lineage assignment changes* in Pangolin



Data: datVis • Chart ID: SankeyID45e8756d4bc8 • googleVis-0.6.3
R version 3.5.2 (2018-12-20) • Google Terms of Use • Documentation and Data Policy

Appendix – VOC Lineages*** Table

VOC	Associated Lineages
Alpha	B.1.1.7, Q.*
Beta	B.1.351, B.1.351.*
Gamma	P.1, P.1.*
Delta	B.1.617.2, AY.*
Omicron	B.1.1.529, BA.*

*indicates an additional numerical value (e.g. Q.1).

***Lineage assignments are based on the use of Pangolin, an epidemiological lineage assignment tool (github.com/cov-lineages/pangolin); these may change with time as new SARS-CoV-2 genomic data becomes available.