



BC Centre for Disease Control
An agency of the Provincial Health Services Authority

ISONIAZID

What is Isoniazid?

Isoniazid is an antibiotic used to prevent or treat tuberculosis (TB).

When should I take this?

- Take on an empty stomach (1 hour before or 2 hours after food)
- Take this medicine at the same time everyday. (This will help you remember to take the medicine)

How should I take this?

- It is very important that you take this medicine regularly.
- Take as directed by your doctor.
- You are to take this medicine:
 - One tablet once each day
 - Other: _____

What should I do if I miss a dose?

- If you miss a dose, take that dose as soon as you remember. But if it is almost time for the next day's dose, skip the missed dose.
- **Do not take a double dose.**

Is there anything I should avoid when taking this?

✗ Alcohol

✗ Antacids

Space at least 1 hour apart from this medicine.

Tell your doctor and pharmacist if you are taking any other medicines, including any over-the-counter (non-prescription) products.

What are some possible side effects?

If you experience the following, stop taking this medicine and report these immediately to your doctor:

- Unusual tiredness or weakness
- Fever
- Loss of appetite
- Nausea (urge to vomit)
- Stomach upset or diarrhea
- Dark urine
- Yellowing of skin or eyes
- Rash

Tell your doctor if you experience burning, tingling, or numbness in your hands or feet. This may mean that you require more Vitamin B6.

It is important that you go for all lab tests ordered by your doctor.

Tell your doctor if you are pregnant, plan to become pregnant, or if you wish to breast-feed your baby.

Where should I store this medication?

- Keep away from the reach of children
- Protect from light
- Tablets: Store at room temperature in a cool, dry place.
- Syrup: Store at room temperature or in the refrigerator.

Where do I call if I have any other questions?