

PYRAZINAMIDE

What is Pyrazinamide?

Pyrazinamide is an antibiotic used only to treat tuberculosis (TB).

When should I take this?

- Take this medicine at the same time everyday.
This will help you remember to take the medicine.

How should I take this?

- It is very important that you take this medicine regularly.
- Take as directed by your doctor.
- You are to take this medicine:
 - Three tablets once each day.
 - Other: _____

What should I do if I miss a dose?

- If you miss a dose, take that dose as soon as you remember. But if it is almost time for the next day's dose, skip the missed dose.
- **Do not take a double dose.**

Is there anything I should avoid when taking this?

X Alcohol

Tell your doctor and pharmacist if you are taking any other medicines, including any over-the-counter (non-prescription) products.

What are some possible side effects?

If you experience the following, stop taking this medicine and report these immediately to your doctor or the Vancouver TB Control Clinic at (604) 707-2692:

- Unusual tiredness or weakness
- Fever
- Loss of appetite
- Nausea (urge to vomit)
- Stomach upset or diarrhea
- Dark urine
- Yellowing of skin or eyes
- Pain or swelling of the joints
- Itching or skin rash

It is important that you go for all lab tests ordered by your doctor. Attend the TB Control Clinic as requested.

Tell your doctor if you are pregnant, plan to become pregnant, or if you wish to breast-feed your baby.

Where should I store this medication?

- Keep away from the reach of children.
- Protect from light.
- Tablets: Store at room temperature in a cool, dry place.
- Suspension: Refrigerate. Shake well before use.

Where do I call if I have any other questions?

Vancouver TB Control Clinic: (604) 707-2692