



## Ministry of Health

### BC Chronic Disease and Selected Procedure Case Definitions

<b>Author:</b>	Chronic Disease Information Working Group
<b>Date Created:</b>	June 29, 2015
<b>Last Updated:</b>	April 4, 2019
<b>Version:</b>	v2017
<b>Email:</b>	<a href="mailto:hlth.cdrwg@gov.bc.ca">hlth.cdrwg@gov.bc.ca</a>

## **GLOSSARY**

### **Cumulative Prevalence**

It is also known as Prevalence for Life. Cases meet case definition criteria to qualify in a given year, and are then carried forward to show as a case every year thereafter. This measure is helpful for determining prevalence related to life-long chronic disease, but less useful for conditions that may be episodic (e.g., mood disorders, substance use problem, etc.).

### **Episodic Prevalence**

Cases meet case definition criteria to qualify in a given year, and are then carried forward in all subsequent years with at least one condition-related code including ICD diagnostic codes, procedure codes or drugs in any specific year. For some conditions (e.g., Gout/crystal arthropathies), the period for at least one recurring diagnosis is the last five years.

### **Annual Prevalence**

Cases meet case definition criteria to qualify in each given year. Can be used as a quality measure to understand how it compares to cumulative prevalence or how it behaves over time.

### **Annual Service Utilization Prevalence**

Cases meet case definition criteria in a given year by virtue of one hospital diagnosis (MRDX) or one physician claim with diagnosis (first position only). Cases are not carried forward from year to year. This utilization prevalence measure is useful for describing the number of people or proportion of the population using services directly related to the disease.

### **Annual Prevalence with 365 Days follow-up**

Cases meet case definition criteria to qualify in each given year with complete 365 days follow-up time. Can be used as a quality measure to understand how it compares to cumulative prevalence or how it behaves over time.

## CASE DEFINITION – DEPRESSION

Case Definition Type: Cumulative Prevalence  Episodic Prevalence  Annual Prevalence  Annual Service Utilization Prevalence   
 Annual Prevalence with 365 Days Follow-up

Case Definition: The case definition for Depression requires one hospitalization or two physician visits in one year with ICD code(s) specified below. The case definition applies to persons aged 1 and older.

Signed-off BC Case Definition: Yes  - V2017 No

Algorithm: 1H or 2P in 1Y with ICD code(s) specified below.

Notes: None

Age Restriction: Age 1 +

Data Source(s):

Data Source	ICD Code/Procedure Code/Rx	ICD Code/Procedure Code Position	Diagnosis Type	Hospital Care Level
Hospital ICD-10	F32, F33	First Position Only <input type="checkbox"/> All Positions <input checked="" type="checkbox"/> Others <input type="checkbox"/> N/A <input type="checkbox"/>	M-Most Responsible Diagnosis <input checked="" type="checkbox"/> 1-Pre-Admit Comorbidity <input checked="" type="checkbox"/> 2-Post-Admit Comorbidity <input checked="" type="checkbox"/> 3-Secondary Diagnosis <input checked="" type="checkbox"/> 4-Morphology Code <input checked="" type="checkbox"/> 5-Admitting Diagnosis <input checked="" type="checkbox"/> 6-Proxy Most Responsible Diagnosis <input checked="" type="checkbox"/> 9-External Cause of Injury Code <input checked="" type="checkbox"/> 0-Newborn <input checked="" type="checkbox"/> W,X,Y- Service Transfer Diagnosis <input checked="" type="checkbox"/>	Acute Care <input checked="" type="checkbox"/> Rehabilitation <input checked="" type="checkbox"/> Day Surgery <input checked="" type="checkbox"/>
Hospital ICD-9	296, 311	First Position Only <input type="checkbox"/> All Positions <input checked="" type="checkbox"/> Others <input type="checkbox"/> N/A <input type="checkbox"/>	M-Most Responsible Diagnosis <input checked="" type="checkbox"/> 1-Pre-Admit Comorbidity <input checked="" type="checkbox"/> 2-Post-Admit Comorbidity <input checked="" type="checkbox"/> 3-Secondary Diagnosis <input checked="" type="checkbox"/> 4-Morphology Code <input checked="" type="checkbox"/> 5-Admitting Diagnosis <input checked="" type="checkbox"/> 6-Proxy Most Responsible Diagnosis <input checked="" type="checkbox"/> 9-External Cause of Injury Code <input checked="" type="checkbox"/> 0-Newborn <input checked="" type="checkbox"/> W,X,Y- Service Transfer Diagnosis <input checked="" type="checkbox"/>	Acute Care <input checked="" type="checkbox"/> Rehabilitation <input checked="" type="checkbox"/> Day Surgery <input checked="" type="checkbox"/>

Physician Claims ICD-9	296, 311, 50B (see note)	First Position Only <input type="checkbox"/> All Positions <input checked="" type="checkbox"/> Others <input type="checkbox"/> N/A <input type="checkbox"/>	N/A	N/A
PharmaNet Drug History	N/A	N/A	N/A	N/A

Note: 50B is a BC Specific diagnostic code used by physician billings (MSP). It is used for both anxiety and depression.

Diagnostic Code:

ICD-9/10	Description
F32	Depressive episode
F33	Recurrent depressive disorder
296	Affective psychoses
311	Depressive disorder, not elsewhere classified
MSP DX Code 50B	Anxiety/Depression

Procedure Code: **N/A**

Drug List: **N/A**