



Ministry of Health

BC Chronic Disease and Selected Procedure Case Definitions

Author:	Chronic Disease Information Working Group
Date Created:	June 29, 2015
Last Updated:	April 4, 2019
Version:	v2017
Email:	hlth.cdrwg@gov.bc.ca

GLOSSARY

Cumulative Prevalence

It is also known as Prevalence for Life. Cases meet case definition criteria to qualify in a given year, and are then carried forward to show as a case every year thereafter. This measure is helpful for determining prevalence related to life-long chronic disease, but less useful for conditions that may be episodic (e.g., mood disorders, substance use problem, etc.).

Episodic Prevalence

Cases meet case definition criteria to qualify in a given year, and are then carried forward in all subsequent years with at least one condition-related code including ICD diagnostic codes, procedure codes or drugs in any specific year. For some conditions (e.g., Gout/crystal arthropathies), the period for at least one recurring diagnosis is the last five years.

Annual Prevalence

Cases meet case definition criteria to qualify in each given year. Can be used as a quality measure to understand how it compares to cumulative prevalence or how it behaves over time.

Annual Service Utilization Prevalence

Cases meet case definition criteria in a given year by virtue of one hospital diagnosis (MRDX) or one physician claim with diagnosis (first position only). Cases are not carried forward from year to year. This utilization prevalence measure is useful for describing the number of people or proportion of the population using services directly related to the disease.

Annual Prevalence with 365 Days follow-up

Cases meet case definition criteria to qualify in each given year with complete 365 days follow-up time. Can be used as a quality measure to understand how it compares to cumulative prevalence or how it behaves over time.

CASE DEFINITION – PERCUTANEOUS TRANSLUMINAL CORONARY ANGIOPLASTY

Case Definition Type: Cumulative Prevalence Episodic Prevalence Annual Prevalence Annual Service Utilization Prevalence
 Annual Prevalence with 365 Days Follow-up

Case Definition: The case definition for Percutaneous Transluminal Coronary Angioplasty (PTCA) requires one hospitalization with procedure code(s) specified below. The case definition applies to persons aged 20 and older.

Signed-off BC Case Definition: Yes - V2017 No

Algorithm: 1H with Procedure Code(s) specified below.

Notes: None

Age Restriction: Age 20 +

Data Source(s):

Data Source	ICD Code/Procedure Code/Rx	ICD Code/Procedure Code Position	Diagnosis Type	Hospital Care Level
CCI	1IJ50, 1IJ57G	First Position Only <input type="checkbox"/> All Positions <input checked="" type="checkbox"/> Others <input type="checkbox"/> N/A <input type="checkbox"/>	M-Most Responsible Diagnosis <input checked="" type="checkbox"/> 1-Pre-Admit Comorbidity <input checked="" type="checkbox"/> 2-Post-Admit Comorbidity <input checked="" type="checkbox"/> 3-Secondary Diagnosis <input checked="" type="checkbox"/> 4-Morphology Code <input checked="" type="checkbox"/> 5-Admitting Diagnosis <input checked="" type="checkbox"/> 6-Proxy Most Responsible Diagnosis <input checked="" type="checkbox"/> 9-External Cause of Injury Code <input checked="" type="checkbox"/> 0-Newborn <input checked="" type="checkbox"/> W,X,Y- Service Transfer Diagnosis <input checked="" type="checkbox"/>	Acute Care <input checked="" type="checkbox"/> Rehabilitation <input checked="" type="checkbox"/> Day Surgery <input checked="" type="checkbox"/>
CCP	48.02, 48.03	First Position Only <input type="checkbox"/> All Positions <input checked="" type="checkbox"/> Others <input type="checkbox"/> N/A <input type="checkbox"/>	M-Most Responsible Diagnosis <input checked="" type="checkbox"/> 1-Pre-Admit Comorbidity <input checked="" type="checkbox"/> 2-Post-Admit Comorbidity <input checked="" type="checkbox"/> 3-Secondary Diagnosis <input checked="" type="checkbox"/> 4-Morphology Code <input checked="" type="checkbox"/> 5-Admitting Diagnosis <input checked="" type="checkbox"/> 6-Proxy Most Responsible Diagnosis <input checked="" type="checkbox"/> 9-External Cause of Injury Code <input checked="" type="checkbox"/> 0-Newborn <input checked="" type="checkbox"/> W,X,Y- Service Transfer Diagnosis <input checked="" type="checkbox"/>	Acute Care <input checked="" type="checkbox"/> Rehabilitation <input checked="" type="checkbox"/> Day Surgery <input checked="" type="checkbox"/>

Physician Claims ICD-9	N/A	First Position Only <input type="checkbox"/> All Positions <input type="checkbox"/> Others <input type="checkbox"/> N/A <input checked="" type="checkbox"/>	N/A	N/A
PharmaNet Drug History	N/A	N/A	N/A	N/A

Diagnostic Code: N/A

Procedure Code:

Data Source	Procedure Code	Description
CCI	1IJ50	Dilation, coronary artery
	1IJ57G	Extraction, coronary arteries using percutaneous transluminal arterial approach
CCP	48.02	Percutaneous transluminal coronary angioplasty without thrombolytic agent
	48.03	Percutaneous transluminal coronary angioplasty with thrombolytic agent

Drug List: N/A