



C. CLINICAL INFORMATION

Date of AIDS diagnosis: YYYY/MM/DD	Date of first positive HIV test: YYYY/MM/DD
First Disease Indicative of AIDS:	
<input type="checkbox"/> Bacterial pneumonia, recurrent * <input type="checkbox"/> Candidiasis (bronchi, trachea or lungs) <input type="checkbox"/> Candidiasis (esophageal) * <input type="checkbox"/> Cervical cancer, invasive <input type="checkbox"/> Coccidioidomycosis (disseminated or extrapulmonary) <input type="checkbox"/> Cryptococcosis (extrapulmonary) <input type="checkbox"/> Cryptosporidiosis (chronic intestinal, >1 mo duration) <input type="checkbox"/> Cytomegalovirus disease (other than in liver, spleen or nodes) <input type="checkbox"/> Cytomegalovirus retinitis (with vision loss) * <input type="checkbox"/> Encephalopathy, HIV-related (dementia) <input type="checkbox"/> Herpes simplex (chronic ulcers, >1 mo duration) <input type="checkbox"/> Histoplasmosis (disseminated or extrapulmonary) <input type="checkbox"/> Isoporiasis (chronic intestinal >1 mo duration)	<input type="checkbox"/> Kaposi sarcoma * <input type="checkbox"/> Lymphoma, Burkitt's <input type="checkbox"/> Lymphoma, immunoblastic <input type="checkbox"/> Lymphoma, primary in brain <input type="checkbox"/> M. avium complex* or M. kansasii*, (disseminated or extrapulmonary) <input type="checkbox"/> M. tuberculosis (disseminated or extrapulmonary) <input type="checkbox"/> M. tuberculosis (pulmonary) <input type="checkbox"/> Mycobacterium of other/unidentified species * <input type="checkbox"/> Pneumocystis carinii pneumonia * <input type="checkbox"/> Progressive multifocal leukoencephalopathy <input type="checkbox"/> Taxoplasmosis of brain <input type="checkbox"/> Wasting syndrome due to HIV

Indicator diseases that apply only to pediatric cases (<15 years old)	
<input type="checkbox"/> Bacterial infections, multiple or recurrent	<input type="checkbox"/> Lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia *

D. RISK FACTORS & EXPOSURE INFORMATION

What are the possible routes of HIV transmission for this client? *Choose all that apply*

<input type="checkbox"/> Sexual exposure	<input type="checkbox"/> Shared syringes	<input type="checkbox"/> Shared other drug using equipment (e.g. crack pipes)	<input type="checkbox"/> Sex work, transactional or exchanges sex	<input type="checkbox"/> Mother is known HIV positive (vertical transmission)
<input type="checkbox"/> Received blood or blood products after 1980, <i>Specify country:</i>	<input type="checkbox"/> Exposure to HIV contaminated blood or body fluids or concentrated virus in an occupational setting, <i>Specify occupation:</i>	<input type="checkbox"/> No identified risk	<input type="checkbox"/> Other <i>please specify:</i>	

E. NOTES



F. CASE DEFINITION

Acquired Immune Deficiency Syndrome (AIDS)

AIDS is listed as a reportable disease in the Communicable Disease Regulation (Schedule A) of the Public Health Act.

AIDS Case definition

One or more of the specified indicator diseases listed in section C, and meets the case definition for HIV infection.

HIV Case definition

Adults, adolescents and children \geq 18 months: Detection of HIV antibody by screening test (i.e., ELISA or Point of Care HIV test) followed by positive confirmatory test (i.e., Western blot or nucleic acid amplification test), or detection of HIV nucleic acid (RNA or DNA) or detection of p24 antigen with confirmation by neutralization assay, or isolation of HIV in culture.

Children < 18 months: Detection of HIV DNA by nucleic acid amplification testing (NAAT) on two separate samples collected at different times.

Ethnicity

Information on ethnic status can be invaluable to public health for planning outreach and public health campaigns. This data is used both to help prevent new infections and to guide care and treatment.

White	e.g. Irish, Scottish, English, Portuguese, Italian, Russian
Black	e.g. African, Haitian, Jamaican, Somali, Nigerian
Chinese	e.g. Chinese, Taiwanese
South Asian	e.g. East Indian, Pakistani, Sri Lankan, Punjabi, Bangladeshi
Southeast Asian	e.g. Vietnamese, Cambodian, Indonesian, Laotian
West Asian	e.g. Afghan, Assyrian, and Iranian
Korean	e.g. Korean only
Japanese	e.g. Japanese only
Filipino	e.g. Filipino only
Latin American	e.g. Mexican, Central/South American
Arab	e.g. Egyptian, Iranian, Moroccan, Lebanese, Kuwaiti, and Libyan
Other	e.g. Use this classification when ethnicity is known but does not appear on list or in cases of dual ethnicity
Unknown	e.g. Use this classification if HCP did not indicate ethnicity
Declined to answer	e.g. Use this classification if patient declined to state self identified ethnicity

Endemic country

Countries where prevalence of HIV among adults (age 15-49 years) is 1.0% or greater **and** one of the following:

- 50% or more of HIV cases are attributed to heterosexual transmission;
- a male to female ratio of 2:1 or less among prevalent infections; or
- HIV prevalence greater than or equal to 2% among women receiving prenatal care.

List of HIV endemic countries can be found at http://www.catie.ca/sites/default/files/HIV-Aids_EpiUpdates_Chapter13_EN.pdf