

Trivalent Inactivated Influenza Vaccine (Inactivated Split Virion) (TIIV) FLUVIRAL®

Supplier: GlaxoSmithKline Inc.

INDICATIONS:

- See [2025/26 Seasonal Influenza Vaccine Eligibility](#)
- **Intended for use in eligible individuals 6 months of age or older, including those with contraindications to LAIV.**

The vaccine is not approved for use in those less than 6 months of age.

DOSES AND SCHEDULE:

Children 6 months-8 years of age (inclusive): 1 or 2 doses given as 0.5 mL **IM**.

Children under 9 years of age who have not previously received any seasonal influenza vaccine require **2 doses** given 4 weeks apart. If the child has received 1 or more doses in any previous season, only a single dose is required. For children requiring 2 doses within the season, it is preferable to use the same vaccine for both doses. However, if the vaccine used for the first dose is not available or unknown, any other age-appropriate vaccine may be used for the second dose.

Children and Adolescents 9-17 years of age (inclusive): 1 dose given as 0.5 mL **IM**.

Adults 18 years of age and older: 1 dose given as 0.5 mL **IM**.

ADMINISTRATION:

Discard multi-dose vials 28 days after first entry.

BOOSTER DOSES AND RE-IMMUNIZATION:

Annually.

SEROLOGICAL TESTING:

Serological testing is not recommended before or after immunization.

CONTRAINDICATIONS:

1. History of anaphylactic reaction to a previous dose of any type of influenza vaccine or to any component of FLUVIRAL®.
2. History of Guillain-Barré syndrome (GBS) within 8 weeks of receipt of a previous dose of influenza vaccine without another cause being identified.

PRODUCT COMPONENTS:

Potential allergens: egg proteins, polysorbate 80, thimerosal (50 µg per 0.5 mL dose; 0.01% w/v) (See SPECIAL CONSIDERATIONS).

Other components: phosphate buffered saline, α-tocopheryl hydrogen succinate, sodium deoxycholate, ethanol, formaldehyde, sucrose.

Trivalent Inactivated Influenza Vaccine (Inactivated Split Virion) (TIIV) FLUVIRAL®

Supplier: GlaxoSmithKline Inc.

PRECAUTIONS:

Severe oculo-respiratory syndrome (ORS) after a previous dose of influenza vaccine. ^A

SPECIAL CONSIDERATIONS:

Egg allergic individuals (including those who have experienced anaphylaxis following egg ingestion) can be immunized with inactivated or live attenuated influenza vaccine in any setting attended by immunization service providers who are following standard vaccine administration practice. ^B

ADVERSE EVENTS:

Local: pain, swelling, redness.

Systemic: fatigue, headache, myalgia, arthralgia, fever, chills, nausea, vomiting, diarrhea.

Infants and toddlers may also experience irritability, drowsiness and loss of appetite. Fewer than 1 in 20 people may develop oculo-respiratory syndrome (ORS). Symptoms include red eyes, a cough, and/or sore throat and/or hoarseness.

^A See [Safety Issues Applicable to Influenza Vaccines, 2. Oculo-Respiratory Syndrome \(ORS\)](#).

^B See [Safety Issues Applicable to Influenza Vaccines, 1. Egg Allergic Individuals](#).