

# BC ROUTINE IMMUNIZATION SCHEDULE

## INFANTS & CHILDREN

| VACCINE                                                                                                                                                                                                                                                       | CHILD'S AGE |          |                                                      |           |           |                                          |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|----------|------------------------------------------------------|-----------|-----------|------------------------------------------|
|                                                                                                                                                                                                                                                               | 2 months    | 4 months | 6 months                                             | 12 months | 18 months | Starting at 4 years (kindergarten entry) |
| <b>DTaP-HB-IPV-Hib</b><br>(diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b)<br>1. HealthLinkBC File # <a href="#">105</a><br>2. Product Monograph <a href="#">Infanrix Hexa</a>                                      | ✓           | ✓        | ✓                                                    |           |           |                                          |
| <b>Pneumococcal conjugate ‡</b><br>1. HealthLinkBC File # <a href="#">62c</a><br>2. Product Monograph <a href="#">Prenar 20</a>                                                                                                                               | ✓           | ✓        |                                                      | ✓         |           |                                          |
| <b>Rotavirus</b><br>1. HealthLinkBC File # <a href="#">104a</a><br>2. Product Monograph <a href="#">Rotarix</a>                                                                                                                                               | ✓           | ✓        |                                                      |           |           |                                          |
| <b>Meningococcal C conjugate</b><br>1. HealthLinkBC File # <a href="#">23a</a><br>2. Product Monograph <a href="#">NeisVac-C</a>                                                                                                                              | ✓           |          |                                                      | ✓         |           |                                          |
| <b>MMR</b><br>(measles, mumps, rubella)<br>1. HealthLinkBC File # <a href="#">14a</a><br>2. Product Monograph <a href="#">MMR II</a> / <a href="#">Priorix</a>                                                                                                |             |          |                                                      | ✓         |           |                                          |
| <b>Varicella ¶</b><br>(chickenpox)<br>1. HealthLinkBC File # <a href="#">44b</a><br>2. Product Monograph <a href="#">Varilrix</a> / <a href="#">Varivax III</a>                                                                                               |             |          |                                                      | ✓         |           |                                          |
| <b>DTaP-IPV-Hib</b><br>(diphtheria, tetanus, pertussis, polio, <i>Haemophilus influenzae</i> type b)<br>1. HealthLinkBC File # <a href="#">15b</a><br>2. Product Monograph <a href="#">Infanrix-IPV/Hib</a> / <a href="#">Pentacel</a>                        |             |          |                                                      |           | ✓         |                                          |
| <b>Tdap-IPV</b><br>(tetanus, diphtheria, pertussis, polio)<br>1. HealthLinkBC File # <a href="#">15a</a><br>2. Product Monograph <a href="#">Adacel-Polio</a> / <a href="#">Boostrix-Polio</a>                                                                |             |          |                                                      |           |           | ✓                                        |
| <b>MMRV ¶</b><br>(measles, mumps, rubella, varicella)<br>1. HealthLinkBC File # <a href="#">14e</a><br>2. Product Monograph <a href="#">Priorix-Tetra</a> / <a href="#">ProQuad</a>                                                                           |             |          |                                                      |           |           | ✓                                        |
| <b>Influenza *</b><br>1. HealthLinkBC File # <a href="#">12d</a> and <a href="#">12e</a><br>2. Product Monograph <a href="#">Afluria Tetra</a> / <a href="#">FluLaval Tetra</a> / <a href="#">Flumist Quadrivalent</a> / <a href="#">Fluzone Quadrivalent</a> |             |          | ✓<br>(yearly for children 6 months of age and older) |           |           |                                          |
| <b>Hepatitis A</b><br>(for Indigenous children only)<br>1. HealthLinkBC File # <a href="#">33</a><br>2. Product Monograph <a href="#">Havrix</a> / <a href="#">Vagta</a>                                                                                      |             |          | ✓                                                    |           | ✓         |                                          |

‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. For more information see the BC Immunization Manual, [Part 4 – Biological Products](#).

¶ Children who had chickenpox or shingles disease, confirmed by a lab test, at 1 year of age or older do not need the chickenpox vaccine and should get the MMR vaccine at 4 years of age.

\* A second dose is needed 4 weeks after the first dose if receiving the vaccine for the first time.

Access the [BC Immunization Manual](#)