

Immunization Programs and Vaccine Preventable Diseases Service 655 West 12th Avenue Vancouver, BC V5Z 4R4

Tel 604.707.2548 Fax 604.707.2515 www.bccdc.ca

Date: July 05, 2016 Administrative Circular: 2016:10

ATTN: Medical Health Officers and Branch Offices

Public Health Nursing Administrators and Assistant Administrators

Holders of Communicable Disease Control Manuals

Re: Update to Communicable Disease Control Manual,
Chapter 2 - Immunization Program,
Section IIA-Immunization Schedules & Section VII-Biological Products

Section IIA-Immunization Schedules

2.0 Routine Schedules

2.1 Schedule A & 2.2 Schedule B
 Meningococcal C conjugate (Men-C-C) vaccine removed from Grade 6, and
 meningococcal quadrivalent conjugate (Men-C-ACYW-135) vaccine added to Grade 9.

2.3 Schedule C

Men-C-C scheduling guidelines revised, and Men-C-ACYW-135 added for children born on or after January 1, 2002 and who are in Grade 9 or older. The accompanying Personalized Schedule Worksheet has been revised to include Men-C-ACYW-135.

Please remove page numbers: 4-7 dated January 2016 Please add new page numbers: 4-7 dated July 2016

3.1 Minimum Intervals Between Vaccine Doses Table

- Meningococcal quadrivalent conjugate vaccines Menveo® and Menactra® added to the table, with associated footnote 'l' relating to the recommended interval between any meningococcal C conjugate vaccine and meningococcal quadrivalent conjugate vaccine being 4 weeks (regardless of which vaccine is given first).
- Footnote 'H' for Men-C-C vaccine revised and content added indicating that NEISVAC-C® is the recommended and supplied product for the primary infant series, however any meningococcal C conjugate vaccine may be used for doses given at 18 months of age or older.

Please remove page numbers: 12 & 13 dated January 2016 Please add new page numbers: 12 & 13 dated July 2016





Section VII-Biological Products

Meningococcal C Conjugate (MCC) Vaccine

- INDICATIONS updated to remove Grade 6 students, and the following eligibility revised to indicate:
 - Children who have not received a dose of Men-C-C vaccine after 12 months of age, and who are younger than Grade 9.
 - Adolescents and adults born prior to January 1, 2002, who have not received a dose
 of Men-C-C vaccine at 10 years of age or older. (Note: a footnote for this indication
 includes a maximum age of 24 years to be eligible for the vaccine.)
- ADMINISTRATION updated to remove content related to Menjugate® Liquid, as BC will
 not be supplying this product.
- SPECIAL CONSIDERATIONS revised to indicate that NEISVAC-C® is the
 recommended and supplied product for the primary infant series, however any
 meningococcal C conjugate vaccine may be used for doses given at 18 months of age or
 older. Content related to the Grade 6 Men-C-C program has also been removed.

Please remove page numbers: 39 & 40 dated January 2016 Please add new page numbers: 39 & 40 dated July 2016

Meningococcal Quadrivalent Conjugate Vaccines (Groups A, C, Y, W-135)

- INDICATIONS updated to include 'adolescents born on or after January 1, 2002 and who
 are in grade 9 or older', with accompanying footnotes for more information. An additional
 footnote has also been added indicating that for medically high risk individuals, Men-CACYW-135 should be given in place of Men-C-C as part of the routine schedule and
 administered according to age at presentation.
- SPECIAL CONSIDERATIONS revisions include:
 - Removal of content indicating that 'these products can be administered to people
 with a history of Guillain-Barre syndrome (GBS) without concern', as this content
 is no longer new and information regarding GBS as a contraindication to a
 particular vaccine can be found in the respective vaccine product pages in
 Section VII-Biological Products, as well as Section IIB-Contraindications and
 Precautions for Immunization.
 - Removal of content indicating that 'high risk children less than 10 years of age
 who have only received one dose of Men-C-ACYW-135 should also receive one
 dose of Men-C-C vaccine if they have not received a dose of Men-C-C vaccine
 after 12 months of age'.

Please remove page numbers: 41 & 42 dated August 2015 Please add new page numbers: 41 & 42 dated July 2016







BC Centre for Disease Control An agency of the Provincial Health Services Authority

If you have any questions or concerns, please contact Christine Halpert, Senior Practice Leader, BCCDC at telephone (604) 707-2555, fax (604) 707-2515 or by email at christine.halpert@bccdc.ca

Sincerely,

Monika Naus, MD MHSc FRCPC FACPM

Medical Director

Immunization Programs and Vaccine Preventable Diseases Service

BC Centre for Disease Control

with hans

pc: BC Ministry of Health:

Dr. Perry Kendall

Provincial Health Officer

Dr. Bonnie Henry

Deputy Provincial Health Officer

Craig Thompson

Director of Immunization

Healthy Populations and Development Branch

Warren O'Briain

Executive Director

Healthy Populations and Development Branch



