BC Centre for Disease Control Provincial Health Services Authority

Referral Form for MMR Vaccination

	Date:		
		YYYY/MM/DD	
MMD VACCINATION OF IMMUNOCOMPROMISED C	LIENTO	DECLUDES DUVSICIAN OD NUDSE	Ī

				Dai					
							YYYY/MM/DI	D	
PRACT familia	TITION r with th	NATION OF IMN ER APPROVAL ne client's curren	: the primary ca	are physician, r		-			
CLIEN	T INFO	RMATION							
Name:									
			Last		First				
DOB:				PHI	N:				
		YYY	Y/MM/DD						
				<u> </u>	II.				
MMR va	accine is	s available for sus	ceptible ^A immur	ocompromised o	clients	s listed belov	W.		
Check	the app	ropriate box for	your client:						
imn	nunosup	ART therapy recipopression has bee ecialist).), inocompetent by a	
☐ Ped	diatric or	ncology treatment	, including autolo	gous HSCT (12	mont	hs after disc	ontinuation of the	rapy).	
□ Acute lymphocytic leukemia in remission for at least 12 months (total lymphocyte count must be ≥1.2 x 10 ⁹ /L), and client not receiving radiation therapy at the time of immunization. If client is still receiving maintenance chemotherapy, it should be withheld for at least 1 week before to 1 week after immunization.									
☐ Chr									
☐ Soli									
☐ Asp	olenia/H	yposplenia (conge	enital, surgical re	moval or functior	nal)				
□ Adults who are no longer immunocompromised due to malignant disease and ≥3 months after completion of immunosuppressive treatment [≥6 months if treated with anti-B-cell antibodies (e.g., rituximab)], not including SOT or HSCT recipients.									
□ ≥1 ı	□ ≥1 month after completion of high dose (≥2 mg/kg or ≥20 mg daily) oral corticosteroid therapy ≥14 days duration.								
□ ≥3 months after completion of immunosuppressive therapy [≥6 months if treated with anti-B-cell antibodies (e.g., rituximab)].									
☐ Isol	☐ Isolated immunodeficiencies: Humoral (Ig), Neutrophil, or complement deficiency diseases								
☐ Oth	☐ Other (specify):								
□ HIV	/-infecte	d client, by age gr	oup ^B						
Immuno	Immunologic <12 months		1-5 years		≥6 years				
category	/	CD4+ T-	Percent (%) of	CD4+ T-		rcent (%) of	CD4+ T-	Percent (%) of	
		lymphocyte	total	lymphocyte	tota		lymphocyte	total lymphocytes	
1		counts (x10 ⁶ /L) ≥1,500	lymphocytes ≥34	counts (x10 ⁶ /L) ≥1000	lyn ≥30	nphocytes	counts (x10 ⁶ /L) ≥500	≥26	
2		750-1,499	26-33	500-999		<u>.</u> -29	200-499	14-25	
								, -	

Individuals with specific health conditions should be immunized with MMR vaccine according to principles outlined in the BC Communicable Disease Control Manual, Chapter 2: Immunization, Part 2 - Immunization of Special Populations.

Communicable Disease Control Manual Chapter 2: Immunization Part 2 – Immunization of Special Populations

A HSCT and CART therapy recipients are considered susceptible regardless of disease or immunization history.

^B NACI recommends that HIV infected individuals who are not severely immunosuppressed (i.e., immunological categories 1 and 2) may be immunized with 2 doses of MMR vaccine separated by 3 months.

Referral Form for MMR Vaccination

					OR LIVE VACCINES: the primary ne client's current medical status.	
	INFORMATION					
Name:						
	L	ast			First	
DOB:			PHN:			
	YYYY	MM/DD				
	To be completed by p	-	urse practitioner enewed after 4 r		to Public Health Nurse.	
I have verified on (YYYY/MM/DD), this client has no medical contraindications to the receipt of live attenuated MMR vaccine. I understand that individuals may require up to 2 doses given up to 3 months apart, and verify that this client's condition is sufficiently stable to permit receipt of 2 doses, during a period of 4 months from the date above.						
Signatur			Clinic:			
Phone #	:		Fax #:			
	•	blic Health Nu	rse and returned	d to physic	cian or nurse practitioner.	
	ealth Nurse Name:					
Phone #			Fax #:			
MMR Va	accine (2 doses, if indica	ated):				
Date:		Lot #:		Site:		
Date:		Lot #:		Site:		