

# BC ROUTINE IMMUNIZATION SCHEDULE INFANTS & CHILDREN

VACCINE	CHILD'S AGE					
	2 months	4 months	6 months	12 months	18 months	Starting at 4 years (kindergarten entry)
<b>DTaP-HB-IPV-Hib</b> (diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # <a href="#">105</a> 2. Product Monograph <a href="#">Infanrix Hexa</a>	✓	✓	✓			
<b>Pneumococcal conjugate ‡</b> 1. HealthLinkBC File # <a href="#">62a</a> 2. Product Monograph <a href="#">Prevnar 13</a>	✓	✓		✓		
<b>Rotavirus</b> 1. HealthLinkBC File # <a href="#">104</a> 2. Product Monograph <a href="#">Rotarix</a>	✓	✓				
<b>Meningococcal conjugate C</b> 1. HealthLinkBC File # <a href="#">23a</a> 2. Product Monograph <a href="#">NeisVac-C</a>	✓			✓		
<b>MMR</b> (measles, mumps, rubella) 1. HealthLinkBC File # <a href="#">14a</a> 2. Product Monograph <a href="#">Priorix</a> / <a href="#">MMR II</a>				✓		
<b>Varicella ¶</b> (chickenpox) 1. HealthLinkBC File # <a href="#">44b</a> 2. Product Monograph <a href="#">Varilrix</a> / <a href="#">Varivax III</a>				✓		
<b>DTaP-IPV-Hib</b> (diphtheria, tetanus, pertussis, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # <a href="#">15b</a> 2. Product Monograph <a href="#">Infanrix-IPV/HIB</a> / <a href="#">Pediacef</a>					✓	
<b>DTaP-IPV</b> (diphtheria, tetanus, pertussis, polio) 1. HealthLinkBC File # <a href="#">15d</a> 2. Product Monograph <a href="#">Infanrix-IPV</a> / <a href="#">Quadracel</a>						✓
<b>MMRV ¶</b> (measles, mumps, rubella, varicella) 1. HealthLinkBC File # <a href="#">14e</a> 2. Product Monograph <a href="#">Priorix-Tetra</a> / <a href="#">ProQuad</a>						✓
<b>Influenza</b> 1. HealthLinkBC File # <a href="#">12d</a> and <a href="#">12e</a> 2. Product Monograph <a href="#">Aqriflu</a> / <a href="#">FluLaval Tetra</a> / <a href="#">Flumist Quadrivalent</a> / <a href="#">Fluviral</a>					✓*	
					(6 months-4 years)	
<b>Hepatitis A ♦</b> 1. HealthLinkBC File # <a href="#">33</a> 2. Product Monograph <a href="#">Avaxim-Pediatric</a> / <a href="#">Havrix</a> / <a href="#">Vaqta</a>			✓		✓	✓ □

- ‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. See HealthLinkBC File # [62a](#) for more information or speak to your health care provider.
- ¶ Children who had chickenpox or shingles disease, diagnosed by a health care provider, at 1 year of age or older do not need the chickenpox vaccine. Children entering school who need both a 2<sup>nd</sup> dose of MMR and varicella vaccines may be immunized with MMRV vaccine.
- \* A second dose is needed 4 weeks after the first dose if receiving vaccine for the first time.
- ♦ Hepatitis A vaccine is offered to Aboriginal children living both on-reserve and off-reserve.
- Hepatitis A vaccine is offered to Aboriginal children starting at 6 months of age. Alternatively, 1 or 2 doses will be offered to kindergarten-age children starting at 4 years of age depending on previous immunization history.

Access the [BC Immunization Manual](#)