

BC ROUTINE IMMUNIZATION SCHEDULE INFANTS & CHILDREN

VACCINE	CHILD'S AGE					
	2 months	4 months	6 months	12 months	18 months	Starting at 4 years (kindergarten entry)
DTaP-HB-IPV-Hib (diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # 105 2. Product Monograph Infanrix Hexa	✓	✓	✓			
Pneumococcal conjugate ‡ 1. HealthLinkBC File # 62a 2. Product Monograph Prevnar 13	✓	✓		✓		
Rotavirus 1. HealthLinkBC File # 104 2. Product Monograph Rotarix	✓	✓				
Meningococcal C conjugate 1. HealthLinkBC File # 23a 2. Product Monograph NeisVac-C	✓			✓		
MMR (measles, mumps, rubella) 1. HealthLinkBC File # 14a 2. Product Monograph Priorix / MMR II				✓		
Varicella ¶ (chickenpox) 1. HealthLinkBC File # 44b 2. Product Monograph Varilrix / Varivax III				✓		
DTaP-IPV-Hib (diphtheria, tetanus, pertussis, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # 15b 2. Product Monograph Infanrix-IPV/HIB / Pediacef					✓	
Tdap-IPV (tetanus, diphtheria, pertussis, polio) 1. HealthLinkBC File # 15a 2. Product Monograph Adacel-Polio / Boostrix-Polio						✓
MMRV ¶ (measles, mumps, rubella, varicella) 1. HealthLinkBC File # 14e 2. Product Monograph Priorix-Tetra / ProQuad						✓
Influenza 1. HealthLinkBC File # 12d and 12e 2. Product Monograph Agriflu / FluLaval Tetra / Flumist Quadrivalent / Fluviral / Fluzone Quadrivalent					✓*	
						(annually for children 6 months to 4 years of age)
Hepatitis A ♦ (for Aboriginal children only) 1. HealthLinkBC File # 33 2. Product Monograph Avaxim-Pediatric / Havrix / Vaqta			✓		✓	

‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. See HealthLinkBC File # [62a](#) for more information or speak to your health care provider.

¶ Children who had chickenpox or shingles disease, diagnosed by a health care provider, at 1 year of age or older do not need the chickenpox vaccine and should get the MMR vaccine at 4 years of age.

* A second dose is needed 4 weeks after the first dose if receiving vaccine for the first time.

♦ Hepatitis A vaccine is offered to Aboriginal children living both on-reserve and off-reserve.

Access the [BC Immunization Manual](#)