## Minimum Intervals Between Eligible Doses: 7-year Olds with Up-to-date Immunizations

Immunization coverage analyses in British Columbia are conducted using business rules which calculate ages and time intervals at receipt of immunization. Each dose is counted as a valid dose only if given at or after the earliest eligible age, or at a time interval equal to or greater than the shortest recommended interval.

| Antigen/Agent   | Minimum Age or Minimum Time Interval<br>Between Eligible Doses |         |                        |          |
|---|--|---------|------------------------|----------|
|   | Dose 1 A   | Dose 2  | Dose 3                 | Dose 4   |
| Diphtheria, Tetanus, acellular Pertussis (DTaP or Tdap) | 42 days  | 28 days | 28 days                | 24 weeks |
| Polio <sup>B</sup>                                      | 42 days  | 28 days | 24 weeks               |          |
| Hepatitis B   |  |         |                        |          |
| received 3rd dose before June 2007                      | 0 days   | 28 days | 28 days                |          |
| received 3rd dose between June 2007 and May 2014        | 0 days   | 28 days | 56 days <sup>c</sup>   |          |
| received 3rd dose in June 2014 or later                 | 0 days   | 28 days | 56 days <sup>C,D</sup> |          |
| Measles   | 12 months  | 28 days |                        |          |
| Mumps   | 12 months  | 28 days |                        |          |
| Rubella <sup>E</sup>                                    | 12 months  |         |                        |          |
| Varicella <sup>F</sup>                                  | 12 months  | 28 days |                        |          |
| Meningococcal C Conjugate <sup>G</sup>                  | 12 months  |         |                        |          |

- A. Dose 1 refers to the earliest age a child can receive the initial dose.
- B. Schedule for DTaP should be followed when poliomyelitis provided in combination vaccine.
- C. Dose 3 must be given at least 16 weeks (112 days) after dose 1.
- D. Dose 3 must be given on or after 24 weeks of age.
- E. Schedule for measles/mumps should be followed when rubella provided in combination vaccine with measles/mumps.
- F. To be counted as valid, varicella vaccine must be administered on or after 12 months of age. Guidelines also state that children with a history of varicella disease should only be considered protected if the illness occurred on or after 12 months of age. The date of varicella disease onset is not systematically entered into iPHIS/Panorama. For the purposes of this assessment, any child with a past history if varicella disease recorded in iPHIS/Panorama is considered protected, regardless of their age at the time of illness.
- G. Dose must be given at least 56 days after any previous meningococcal C conjugate dose (if previous dose given).

British Columbia immunization coverage reports and definitions are available at: <a href="http://www.bccdc.ca/health-info/immunization-vaccines/immunization-coverage">http://www.bccdc.ca/health-info/immunization-vaccines/immunization-coverage</a>.