



BC Centre for Disease Control
AN AGENCY OF THE PROVINCIAL HEALTH SERVICES AUTHORITY

STI

Annual Report
2015

Contact Information

BC Centre for Disease Control
Clinical Prevention Services
655 West 12th Avenue
Vancouver BC V5Z 4R4
Phone: 604-707-5621
Fax: 606-707-5604
Email: CPSSurveillance@bccdc.ca

Date of publication: August 31, 2017
Report is available at www.bccdc.ca

Suggested citation:

BC Centre for Disease Control. (2017). STI in British Columbia: Annual Surveillance Report 2015.
Retrieved from <http://www.bccdc.ca/search?k=sti%20annual%20report>



Table of Contents

Summary of Trends	4
Chlamydia	5
Genital Chlamydia by Region, Gender, and Age	5
Extra-genital Chlamydia	10
Perinatally-acquired Chlamydia	11
Lymphogranuloma Venereum	12
Gonorrhea	13
Genital Gonorrhea by Region, Gender, and Age	13
Extra-genital Gonorrhea	19
Perinatally-acquired Gonorrhea	19
Gonorrhea Antimicrobial Resistance	20
Pelvic Inflammatory Disease and Ectopic Pregnancy	22
Infectious Syphilis	24
Infectious Syphilis by Region, Gender, and Age	24
Infectious Syphilis by Ethnicity	29
Infectious Syphilis among Aboriginal Peoples	30
Infectious Syphilis by Exposure Category	31
Infectious Syphilis among Men who have Sex with Men	34
Stage of Infection at Time of Syphilis Diagnosis	36
Maternal and Early Congenital Syphilis	37
Endnotes	38
Contributors	40
Technical Appendix	41
Data Limitations	41
Case Definitions	42
Data Sources	43
Additional Notes	45

Summary of Trends

Genital Chlamydia

In 2015, the rate of genital chlamydia increased to 304.0 per 100,000 population, continuing the overall provincial trend of a steady increase since 1998.

- The highest rates were in Vancouver, Northwest, and Northern Interior Health Service Delivery Areas.
- Females continue to have higher rates of genital chlamydia infection compared to males.
- The highest rates were among young adults aged 20-29 years followed by adolescents aged 15-19 years.
- In 2015, there were 166 extra-genital infections identified and 3 perinatally-acquired infections.
- The number of LGV cases remains higher than historic levels. In 2015, the number of LGV cases reached an all-time high of 42 cases, all were among men who have sex with men, and many of whom are co-infected with HIV.

Genital Gonorrhea

In 2015, the provincial rate of genital gonorrhea increased to 67.4 per 100,000 population, continuing an overall steady increase since 1998.

- The highest rates were in Vancouver, Northwest, and Fraser East Health Service Delivery Areas.
- Males have higher rates of infection compared to females and in 2015, rates among both males and females increased to an all-time high.
- The highest rates of infection were among males aged 25-29 years and among females aged 20-24 years.
- In 2015, there were 505 extra-genital infections identified and one perinatally-acquired infection.
- Analysis of recent gonorrhea antimicrobial resistance trends in BC demonstrates a reduction in the proportion of isolates with reduced susceptibility to cefixime and ceftriaxone since 2011.

Pelvic Inflammatory Disease and Ectopic Pregnancy

In 2015, the rate of physician billings and hospital admissions for women related to pelvic inflammatory disease was stable or decreased which is consistent with the overall decline in this potential complication of chlamydia or gonorrhea infection over the past two decades. While hospitalizations of women related to ectopic pregnancy have shown a similar trend, the number of physician billings has shown a small but steady increase since 2003 in BC.

Infectious Syphilis

In 2015, the provincial rate of infectious syphilis increased to 16.2 per 100,000 population, continuing a steady increase since 2011.

- The highest rates of infection were in Vancouver and South Vancouver Island Health Service Delivery Areas.
- In 2015, over 95% of cases were male, with highest rates observed in males aged 25-29 years. A slight increase in female cases was identified in 2015.
- The majority of cases in 2015 were among people identified as Caucasian (48%).
- Men who have sex with men (MSM) continue to comprise the greatest number of new infectious syphilis cases in BC (84% in 2015). Among MSM cases where HIV status is known, 51% were co-infected with HIV.

Chlamydia

Genital Chlamydia by Region, Gender, and Age


Genital chlamydia is the most commonly reported sexually transmitted infection in BC. As the majority of chlamydia infections are asymptomatic, the reported number of chlamydia infections is only a portion of the total population burden. If untreated, genital chlamydia may lead to complications such as pelvic inflammatory disease (a major cause of infertility, ectopic pregnancy, and chronic pain) in women and epididymo-orchitis in men.

Mirroring the national trend, genital chlamydia rates have steadily increased from 1998 to 2015 following a decline in the early 1990s (Figure 2). In 2015, the rate of genital chlamydia for BC increased to 304.0 (14,245 cases) from 288.3 (13,352 cases) per 100,000 population in 2014. The highest rates of genital chlamydia were in Vancouver Coastal and Northern Health Authorities (Figure 3). Rates among Health Service Delivery Areas varied with the highest rates in Vancouver, Northwest and Northern Interior, and the lowest rates in Kootenay Boundary, Fraser East and Fraser South (Figure 1).

Similar increases in chlamydia infections have been observed in high income countries around the world.^{1,2} There are multiple reasons for this increase, including increases in the sensitivity of laboratory tests and uptake of testing (e.g., the greater acceptability of urine-based tests among men) as well as provider screening practices. There may also be a paradoxical effect in which improvements in early screening and treatment for chlamydia over the past decades have resulted in individuals being less likely to develop full immunity thus consequently more susceptible to re-infection (known as the “arrested immunity” hypothesis^{3,4}). While data on population trends in sexual behaviour is not available for BC, it is possible that changes in behaviour, such as decreased condom use, may also be contributing to increasing chlamydia incidence.


Females continue to have approximately 1.6 times the diagnosis rate compared to males. The rate for females in 2015 increased to 369.7 (8,724 cases) from 352.8 (8,225 cases) per 100,000 population in 2014 (Figure 4). The greater number of infections among females is partially due to routine screening performed at the time of visits that were for other reasons (e.g., pap testing or contraception counselling). In 2015, the highest rates of chlamydia were among young adults aged 20-29 years followed by adolescents aged 15-19 years (Figure 6), driven primarily by the high rates of infection among young females. For females, the rates for all age groups increased from 2014 to 2015 (Figure 7). Males aged 20-29 years had the highest chlamydia rates in 2015 compared with other age groups (Figure 8).

1. Genital chlamydia case reports in BC by health service delivery area, 2015




Rates calculated with population estimates released by BC Stats


2. Genital and extra-genital chlamydia case reports in BC and Canada, 1991 to 2015



3. Genital chlamydia case reports in BC by health authority, 2006 to 2015




4. Genital chlamydia case reports in BC by gender, 2006 to 2015




* Other - transgender and gender unknown

5. Genital chlamydia case reports in BC by age group and gender, 2015




* Other - transgender and gender unknown


6. Genital chlamydia case reports in BC by age group - total, 2006 to 2015



7. Genital chlamydia case reports in BC by age group - female, 2006 to 2015



8. Genital chlamydia case reports in BC by age group - male, 2006 to 2015



Extra-genital Chlamydia

In 2015, 166 extra-genital cases were identified (40 females, 124 males) which is an increase from 116 cases in 2014 (25 females and 91 males). As screening for chlamydia infections at extra-genital sites is not routine practice, these findings are strongly influenced by provider testing practices. Much of the increase observed since 2012 is likely due to increase awareness and testing for chlamydia in extra-genital sites. From 2006 to 2015, 514 infections were identified in specimens collected from the following sites: throat (360 cases, 70.0%), eye (95 cases, 18.5%), lung (1 case, 0.2%), and other sites (58 cases, 11.3%) (Table 9).

9. Extra-genital chlamydia case reports in BC by site/culture, 2006 to 2015

Gender	Site	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Female	Throat	1	2	0	0	0	0	6	5	9	30
	Eye	6	3	3	6	3	3	2	2	6	4
	Lung	0	0	0	0	0	0	0	0	0	0
	Other *	0	1	1	5	5	4	1	4	10	6
	Total	7	6	4	11	8	7	9	11	25	40
Male	Throat	4	1	6	2	7	1	51	42	80	109
	Eye	8	5	5	5	5	5	3	2	8	11
	Lung	0	1	0	0	0	0	0	0	0	0
	Other *	0	0	2	0	3	5	2	2	3	4
	Total	12	7	13	7	15	11	56	46	91	124
BC	Throat	5	3	6	2	7	1	59	47	89	141
	Eye	14	8	8	11	8	8	5	4	14	15
	Lung	0	1	0	0	0	0	0	0	0	0
	Other *	0	1	3	5	8	9	3	6	13	10
	Total	19	13	17	18	23	18	67	57	116	166


* Other - nasopharyngeal washing, lesion, fluid from groin mass lesion, pelvic fluid, and other

Total reports for BC is the sum of the following genders: female, male, transgender, and gender unknown

Perinately-acquired Chlamydia

Genital chlamydia can be passed by pregnant women to their infants during delivery which can lead to ophthalmia neonatorum and pneumonia. Three cases of perinately-acquired chlamydia infection were found in conjunctival specimens in 2015. From 2006 to 2015, the majority of perinatal cases are from conjunctival specimens (18/19 cases, 94.7%) while one case (5.3%) was identified in a respiratory specimen (Figure 10).

10. Perinately-acquired chlamydia case reports in BC by site, 2006 to 2015



Lymphogranuloma Venereum

Lymphogranuloma venereum (LGV) is a sexually transmitted infection caused by *Chlamydia trachomatis* serovars L1, L2, and L3. The clinical presentation of LGV includes genital papules, ulcers, inguinal lymphadenopathy, and hemorrhagic proctitis. If left untreated, LGV can cause serious sequelae such as lymphatic obstruction or anogenital ulcerations. LGV can easily be misdiagnosed as other sexually transmitted infections or gastrointestinal disease.

LGV was first reported in Canada in 2003 and in BC in 2004. With increasing cases of LGV among gay, bisexual, and other men who have sex with men (MSM) in Europe and the US, provincial LGV surveillance commenced in 2004. Occurring in tandem with reports of increased transmission in the US⁵ and Europe⁶, an increase of LGV cases was observed in 2011 in BC, in part due to routine testing of rectal chlamydia specimens for LGV and augmented case-finding. In 2015, the number of LGV cases in BC reached an all-time high of 42 cases (40 confirmed, 2 probable) (Figure 11).

From 2006 to 2015, 141 cases of LGV (122 confirmed, 19 probable) were reported in BC. Most cases (140 cases, 99.3%) were among MSM and diagnosed in Vancouver (110 cases, 78.0%). Of those with known HIV status, 67.9% (91/134 cases) were co-infected with HIV. Most cases (110/123 cases, 89.4%) presented with symptoms of proctitis. In 2015, the rate of LGV in BC was 1.8 (42 cases) per 100,000 population among males; all cases were MSM. The mean age was 41.2 years (range 22-62 years), and 61.9% identified as Caucasian (26 cases).^{7, 8}

11. Lymphogranuloma venereum case reports in BC, 2006 to 2015



Gonorrhea

Genital Gonorrhea by Region, Gender, and Age

Gonorrhea infections may be asymptomatic or the symptoms may be mild, and as a result, the reports of diagnosed infections are only a portion of the total population burden. As with chlamydia, genital gonorrhea if untreated can lead to pelvic inflammatory disease (and associated complications) in females. An individual infected with gonorrhea is also at increased risk of acquiring HIV.^{9, 10}


Overall since 1998, the provincial genital gonorrhea rate in BC has increased steadily, consistent with national rates (Figure 13). However, the genital gonorrhea rate in BC increased over 70% to 67.4 (3,160 cases) in 2015 from 38.9 (1,803 cases) per 100,000 population in 2014. The highest rate was in the Vancouver Coastal Health Authority (Figure 14). Rates among Health Service Delivery Areas vary with the highest rates in Vancouver, Northwest and Fraser East, and the lowest rates in East Kootenay, Richmond and North Vancouver Island (Figure 12). Reasons for the increase in gonorrhea rates in 2015 are still under investigation. Preliminary findings suggest that a shift in strain-type may account for some of this increase, as well as increased testing. The BCCDC is working in partnership with the BCCDC Public Health Laboratory (BCCDC PHL), National Microbiology Laboratory (NML) and Public Health Agency of Canada (PHAC) to understand the epidemiology of gonorrhea.

Males continue to have a rate about two times greater than females (Figure 15). Since 2006, male and female gonorrhea rates have been steadily increasing. In 2015, the rates among both males and females increased to a 25-year high of 88.8 (2,065 cases) among males and 46.2 (1,089 cases) among females per 100,000 population.

Similar to trends from 2006 to 2014, in 2015, the highest rates of gonorrhea were among those aged 20-29 years (Figure 17). In 2015, the highest rates among males were in those aged 25-29 years (445 cases, 280.7 per 100,000 population) and among females in those aged 20-24 years (301 cases, 194.8 per 100,000 population) (Figure 16).


Gonorrhea is more likely to be concentrated in sexually active networks and it is likely that the higher rates of gonorrhea in males is, in part, due to higher rates of gonorrhea among gay, bisexual, and other men who have sex with men (MSM). While provincial surveillance data does not permit identification of MSM cases, this has been observed in other jurisdictions.^{11, 12}

12. Genital gonorrhea case reports in BC by health service delivery area, 2015




Rates calculated with population estimates released by BC Stats


13. Genital and extra-genital gonorrhea case reports in BC and Canada, 1991 to 2015



14. Genital gonorrhea case reports in BC by health authority, 2006 to 2015




15. Genital gonorrhea case reports in BC by gender, 2006 to 2015




* Other - transgender and gender unknown

16. Genital gonorrhea case reports in BC by age group and gender, 2015




* Other - transgender and gender unknown


17. Genital gonorrhea case reports in BC by age group - total, 2006 to 2015



18. Genital gonorrhea case reports in BC by age group - female, 2006 to 2015



19. Genital gonorrhea case reports in BC by age group - male, 2006 to 2015



Extra-genital Gonorrhea

In 2015, 505 cases were identified (41 females, 464 males) which was an increase from 366 cases (22 females, 344 males) in 2014. As screening for gonorrhea infections at extra-genital sites is not routine practice, these findings are strongly influenced by provider testing practices, including increases in screening in the past few years. Of the 1,678 cases diagnosed from 2006 to 2015, cases were identified in the throat (1,608 cases, 95.8%), eye (35 cases, 2.1%), and other sites (29 cases, 1.7%). A small number of the diagnosed cases represented disseminated gonococcal infection (6 cases, 0.4%) (Table 20).

20. Extra-genital gonorrhea case reports in BC by site/culture, 2006 to 2015

Gender	Site	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Female	Throat	16	15	3	7	8	13	8	26	19	37
	Eye	0	1	1	0	1	0	0	1	1	4
	Other *	0	4	0	0	1	0	0	1	2	0
	DGI **	0	0	1	0	0	0	1	1	0	0
	Total	16	20	5	7	10	13	9	29	22	41
Male	Throat	41	46	41	44	55	95	155	184	334	454
	Eye	0	1	1	5	4	0	1	1	6	7
	Other *	1	0	0	1	4	5	2	1	4	3
	DGI **	0	0	1	1	0	0	1	0	0	0
	Total	42	47	43	51	63	100	159	186	344	464
BC	Throat	57	61	44	53	63	108	165	213	353	491
	Eye	0	2	2	5	5	0	1	2	7	11
	Other *	1	4	0	1	5	5	2	2	6	3
	DGI **	0	0	2	1	0	0	2	1	0	0
	Total	58	67	48	60	73	113	170	218	366	505

* Other - superficial wound, sternoclavicular synovium, synovium joint fluid, abscess, blood, and elbow

** DGI - disseminated gonococcal infection

Total reports for BC is the sum of the following genders: female, male, transgender, and gender unknown

Perinatally-acquired Gonorrhea

In 2015, there was one report of perinatally-acquired gonorrhea. From 2006 to 2015, four perinatal cases have been identified.

Gonorrhea Antimicrobial Resistance

Treatment of gonorrhea has long been challenged by the bacterium's ability to acquire resistance to multiple classes of antibiotics. Historically, effective antibiotics – penicillin, doxycycline, and ciprofloxacin – can no longer be used successfully against gonorrhea, leaving few remaining options. BC treatment guidelines currently recommend third-generation cephalosporins for the treatment of gonorrhea: injectable ceftriaxone (250mg) or oral cefixime (800mg), co-treated with 1g of azithromycin.¹³ Recent international surveillance data and case reports however, suggest that susceptibility of gonorrhea to these current first-line treatments is also now threatened.^{14, 15} In this context, local surveillance is critical.

The BCCDC Public Health Laboratory (PHL) routinely tests *N. gonorrhoeae* isolates for susceptibility to a panel of antimicrobial drugs, including cefixime, ceftriaxone, and azithromycin. Data presented here summarize the minimum inhibitory concentration (MIC) of these drugs among isolates from BC. The MIC is the lowest amount of antibiotic required to inhibit growth of the bacterium; a higher MIC means the bacterium is less susceptible to the antibiotic.


A total of 4,180 isolates were tested between 2006 and 2015, representing 24.6% (4,180/16,964) of all gonorrhea cases reported in the province. Fifty percent (2,060/4,180) of isolates tested for drug susceptibility were sampled from the urethra, 25.3% (1,056/4,180) from the rectum, 12.1% (506/4,180) from the cervix, and 12.4% (519/4,180) from the throat.

Since 2006, 0.5% (21/4,180) of isolates showed an MIC ≥ 0.25 µg/mL to cefixime. Fortunately, no isolate was fully resistant to cefixime or ceftriaxone¹⁶ and no treatment failures were reported in BC during this period (Figure 21). The increasing trend in percentage of isolates with elevated MIC (i.e., reduced susceptibility) to cefixime or ceftriaxone observed in 2006-2010 reversed in 2011-2015. Similarly, the increasing trend in percentage of isolates with elevated MIC to azithromycin¹⁷ in 2006-2011 reversed in 2012-2015.

The decline observed in 2011-2015 for reduced susceptibility to cefixime or ceftriaxone among tested isolates is encouraging and, may in part be due to changes in the Canadian and provincial gonorrhea treatment guidelines to more effective regimens (i.e., increased cefixime dosage or improved medication adherence due to single dosage). These trends will be closely monitored in order to inform the future evolution of gonorrhea treatment recommendations. The continued threat of emerging resistance reinforces the need for STI prevention and control measures such as increased testing for gonorrhea, partner testing and treatment of gonorrhea, and tests of cure, as well as the need for antibiotic stewardship to ensure effective treatments for bacterial infections.

21. Percentage of tested *N. gonorrhoeae* isolates with elevated minimum inhibitory concentrations (MIC) to Cefixime, Ceftriaxone, and Azithromycin in BC, 2006 to 2015

Elevated MIC defined here as ≥ 0.064 $\mu\text{g/mL}$ for cefixime/ceftriaxone and ≥ 0.5 $\mu\text{g/mL}$ for azithromycin



Pelvic Inflammatory Disease and Ectopic Pregnancy

Pelvic inflammatory disease (PID) and ectopic pregnancy (EP) are medical conditions in women that can be caused by chlamydia or gonorrhea infection. Examination of the rates of these conditions can provide an indication of the complications of chlamydia and gonorrhea infections. Included in this report are data of physician billings (representing inpatient and outpatient treatment) and hospital discharges (representing inpatient treatment) provided by the BC Ministry of Health.

In BC, steady declines in physician billings and hospital discharges related to PID have been observed over time. Rates in hospital discharges for EP have shown a similar trend although physician billings for EP have increased slightly in recent years. Despite increasing rates of chlamydia and gonorrhea infections, PID has been declining. This paradoxical trend may be a result of targeted screening programs that have led to earlier diagnosis and treatment of chlamydia and gonorrhea that may reduce the probability of complications such as PID and EP.


Pelvic Inflammatory Disease

In 2015, the rate of physician billings related to PID decreased slightly to 221.7 (2,031 physician billings) from 223.8 (2,043 physician billings) per 100,000 women aged 15-44 years in 2014 (Figure 22). Rate of hospital discharges related to PID also show a decrease to 25.1 (230 hospital discharges) in 2015 from 29.8 (272 hospital discharges) per 100,000 women aged 15-44 years in 2014 (Figure 23).


Ectopic Pregnancy

The rate of physician billings related to EP increased to 267.0 (2,446 physician billings) in 2015 from 239.7 (2,188 physician billings) per 100,000 women aged 15-44 years in 2014 (Figure 22). Rate of hospital discharges related to EP have remained the same in 2015 at 50.2 (460 hospital discharges) as in 2014 at 49.4 (451 hospital discharges) per 100,000 women aged 15-44 years (Figure 23).

22. Case reports of women aged 15-44 years with a physician billing related to pelvic inflammatory disease or ectopic pregnancy in BC, 1993 to 2015



23. Case reports of women aged 15-44 years with a hospital discharge related to pelvic inflammatory disease or ectopic pregnancy in BC, 1993 to 2015



Infectious Syphilis

Infectious Syphilis by Region, Gender, and Age


Syphilis infections are divided into several stages: primary, secondary, early latent, and late latent. The initial symptoms of syphilis may not always be recognized and without treatment individuals generally enter a prolonged asymptomatic phase. Individuals can still however, be infectious despite not having any symptoms. Syphilis infection can lead to serious complications, including cardiovascular and neurologic disease, and may be fatal.

Following a decline in rates in BC in the early 1990's, infectious syphilis (i.e., primary, secondary, and early latent stages) began to re-emerge in BC starting in 1997, corresponding to a series of outbreaks in different populations. While provincial trends had been decreasing in 2009-2010, infectious syphilis rates began to increase in 2011 and this became more evident in 2012-2013. In BC, the rate of infectious syphilis increased in 2015 to 16.2 (761 cases) from 11.9 (549 cases) per 100,000 population in 2014 (Figure 24). The highest rate of infectious syphilis was in the Vancouver Coastal Health Authority (Figure 26). Across Health Service Delivery Areas, the highest rates were in Vancouver and South Vancouver Island (Figure 25).


The majority of infectious syphilis cases in BC are male (Figure 27). Although male infectious syphilis rates decreased in 2009-2010, male infectious syphilis rates have since risen to 31.5 (732 cases) per 100,000 population in 2015. The increase in cases among males was observed in all age groups. Since 2014, the highest rates of syphilis was among males 25-29 years old, followed closely by males 30-39 years old (Figure 31). In 2015, the rate of infectious syphilis among females was 1.2 (29 cases), a slight increase from 1.0 (24 cases) per 100,000 population in 2014 (Figure 27).

In 2015, 628 cases (82.5% of all cases) reported sexual (including anonymous) partners while the remaining cases (133 cases, 17.5%) provided no information regarding their sexual partners or were lost to follow-up. Of these 628 cases, 465 cases (74.0%) reported 2,861 sexual partners (notifiable and anonymous); 178 cases (28.3%) reported anonymous partners. Most of the cases (491 cases, 78.2%) stated they would notify their sexual partners themselves.

24. Infectious syphilis case reports in BC and Canada, 1991 to 2015*




25. Infectious syphilis case reports in BC by health service delivery area, 2015




Rates calculated with population estimates released by BC Stats

26. Infectious syphilis case reports in BC by health authority, 2006 to 2015




27. Infectious syphilis case reports in BC by gender, 2006 to 2015




* Other - transgender and gender unknown

28. Infectious syphilis case reports in BC by age group and gender, 2015




* Other - transgender and gender unknown


29. Infectious syphilis case reports in BC by age group - total, 2006 to 2015



30. Infectious syphilis case reports in BC by age group - female, 2006 to 2015



31. Infectious syphilis case reports in BC by age group - male, 2006 to 2015



Infectious Syphilis by Ethnicity

In males, almost half of cases (365 cases; 49.9%) in 2015 were among people who identified as Caucasian (Table 34). In comparison, most females (5 cases, 17.2%) in 2015 identified as Aboriginal however, the trends are highly variable due to the small number of female cases each year (29 cases in 2015) (Table 33).

32. Percentage of infectious syphilis case reports in BC by ethnicity - total, 2006 to 2015

Ethnicity	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
<i>No. Diagnoses</i>	331	301	328	217	154	191	369	558	549	761
Caucasian	60.4	64.5	66.2	65.0	70.1	64.9	61.2	55.7	52.3	48.5
Aboriginal	12.4	10.6	8.8	7.8	3.9	3.7	5.7	3.9	2.9	3.2
Asian	11.5	8.6	10.7	6.5	12.3	9.9	10.6	12.2	13.5	8.9
South Asian	6.0	4.0	4.3	1.4	3.2	6.8	4.1	3.6	2.6	2.2
Hispanic	4.2	5.6	4.9	9.2	2.6	8.4	6.2	5.2	6.0	5.5
Black	2.7	2.3	2.4	4.6	1.3	0.5	1.1	1.6	2.4	0.5
Other*	2.4	2.0	1.2	3.7	2.6	3.1	1.9	1.3	1.5	1.7
Unknown	0.3	2.3	1.5	1.8	3.9	2.6	9.2	16.5	18.9	29.4

* Other - Arab/West Asian and other/mixed ethnicity

33. Percentage of infectious syphilis case reports in BC by ethnicity - female, 2006 to 2015

Ethnicity	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
<i>No. Diagnoses</i>	77	55	41	28	13	13	24	20	24	29
Caucasian	40.3	54.5	43.9	57.1	69.2	38.5	29.2	15.0	20.8	13.8
Aboriginal	42.9	34.5	29.3	28.6	15.4	15.4	20.8	15.0	8.3	17.2
Asian	7.8	5.5	12.2	0.0	0.0	23.1	16.7	30.0	37.5	10.3
South Asian	6.5	3.6	9.8	7.1	7.7	7.7	12.5	10.0	4.2	0.0
Hispanic	0.0	0.0	0.0	0.0	0.0	7.7	0.0	5.0	0.0	3.4
Black	1.3	0.0	2.4	3.6	7.7	0.0	4.2	0.0	0.0	0.0
Other*	1.3	0.0	0.0	3.6	0.0	0.0	0.0	0.0	0.0	3.4
Unknown	0.0	1.8	2.4	0.0	0.0	7.7	16.7	25.0	29.2	51.7

* Other - Arab/West Asian and other/mixed ethnicity

34. Percentage of infectious syphilis case reports in BC by ethnicity - male, 2006 to 2015

Ethnicity	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
<i>No. Diagnoses</i>	254	245	286	188	141	178	344	536	524	732
Caucasian	66.5	66.9	69.6	66.5	70.2	66.9	63.4	57.3	53.8	49.9
Aboriginal	3.1	5.3	5.6	4.8	2.8	2.8	4.7	3.5	2.7	2.6
Asian	12.6	9.4	10.5	7.4	13.5	9.0	10.2	11.6	12.4	8.9
South Asian	5.9	4.1	3.5	0.5	2.8	6.7	3.5	3.4	2.5	2.3
Hispanic	5.5	6.5	5.6	10.6	2.8	8.4	6.7	5.2	6.3	5.6
Black	3.1	2.9	2.4	4.8	0.7	0.6	0.9	1.7	2.5	0.5
Other*	2.8	2.4	1.4	3.2	2.8	3.4	2.0	1.1	1.5	1.6
Unknown	0.4	2.4	1.4	2.1	4.3	2.2	8.7	16.2	18.3	28.6

* Other - Arab/West Asian and other/mixed ethnicity


Infectious Syphilis among Aboriginal Peoples

This section describes cases of infectious syphilis among people who identify as Aboriginal. Among the nearly 200,000 Aboriginal persons living in BC, approximately 66% are First Nations, 30% are Métis, and fewer than 5% are Inuit or of other Aboriginal identity.¹⁸

Since 2006, the percentage of infectious syphilis cases in BC residents that were reported among Aboriginal Peoples has been decreasing. In 2015, the percentage of infectious syphilis cases among Aboriginal Peoples increased slightly to 3.2% (24 cases) from 2.9% (16 cases) in 2014 (Table 32). In comparison, Aboriginal Peoples represent about 5% of the general BC population.¹⁹ While numbers are small, Aboriginal women in BC are disproportionately represented among all female infectious syphilis cases (5 cases or 17.2% of all female cases in 2015).

Consistent with overall provincial trends, the rate of infectious syphilis among First Nations people in BC has increased since 2010 (Figure 35). The rate increased in 2015 to 15.4 (22 cases) from 11.4 (16 cases) per 100,000 population in 2014. Due to limitations in the availability of population estimates of Métis and Inuit peoples, it is not possible to calculate comparable rates. (See Technical Appendix for further details about the classification of ethnicity for syphilis cases and First Nations population estimates.)

35. Infectious syphilis case reports among First Nations people in BC by gender, 2006 to 2015



Rates based on First Nations population estimates from Aboriginal Affairs and Northern Development Canada (AANDC)

Infectious Syphilis by Exposure Category

Gay, bisexual, and other men who have sex with men (MSM) continue to comprise the greatest number of infectious syphilis cases in BC (Figure 37). The number of syphilis cases among MSM increased to 642 cases (84.4% of all cases) in 2015 from 466 cases (84.9%) in 2014. Infectious syphilis cases among heterosexual persons without other risk factors increased in 2015 (82 cases, 10.8% in 2015; 51 cases, 9.3% in 2014). The number of infectious syphilis cases among street involved persons, sex trade workers, and their patrons has continued to decrease (6 cases, 1.1% in 2014; 4 cases, 0.5% in 2015). Trends among MSM in BC are explored in more detail in the next section.

36. Infectious syphilis case reports in BC by exposure category and health authority, 2006 to 2015

Health Authority	Exposure Category	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Interior	MSM	2	7	2	5	1	3	6	4	3	23
	Street/STW	1	3	3	1	1	0	0	0	0	0
	HET	2	0	2	7	2	4	2	2	2	6
	Outside	0	0	0	0	0	0	1	0	1	0
	Other/UNK	0	0	0	0	0	0	0	0	1	0
Fraser	MSM	22	13	41	25	17	15	46	71	78	105
	Street/STW	34	23	20	4	2	2	4	4	1	2
	HET	25	21	15	13	5	8	12	19	11	32
	Outside	3	3	5	1	2	0	1	2	2	0
	Other/UNK	2	1	1	0	0	2	1	2	4	7
Vancouver Coastal	MSM	129	136	179	108	94	121	248	386	356	460
	Street/STW	62	34	23	8	9	6	9	10	5	2
	HET	27	14	14	18	11	7	18	24	29	30
	Outside	6	6	1	2	3	1	0	3	4	1
	Other/UNK	0	1	1	1	1	2	4	8	13	19
Vancouver Island	MSM	4	7	6	7	3	13	10	13	22	42
	Street/STW	0	3	3	4	0	0	0	0	0	0
	HET	8	7	2	7	1	3	2	6	8	12
	Outside	1	0	0	0	0	0	0	0	1	0
	Other/UNK	0	0	1	1	0	0	1	1	0	6
Northern	MSM	0	6	1	0	0	2	0	1	4	10
	Street/STW	2	2	4	2	0	1	1	0	0	0
	HET	1	8	4	1	2	1	3	0	1	1
	Outside	0	0	0	0	0	0	0	0	0	0
	Other/UNK	0	0	0	0	0	0	0	1	0	0
nonBC	MSM	0	5	0	1	0	0	0	1	3	2
	Street/STW	0	0	0	1	0	0	0	0	0	0
	HET	0	0	0	0	0	0	0	0	0	1
	Outside	0	0	0	0	0	0	0	0	0	0
	Other/UNK	0	1	0	0	0	0	0	0	0	0
BC	MSM	157	174	229	146	115	154	310	476	466	642
	Street/STW	99	65	53	20	12	9	14	14	6	4
	HET	63	50	37	46	21	23	37	51	51	82
	Outside	10	9	6	3	5	1	2	5	8	1
	Other/UNK	2	3	3	2	1	4	6	12	18	32

MSM - men who have sex with men


Street/STW - street involved / sex trade worker or patron

HET - heterosexual contact


Outside - acquired outside of Canada

UNK - exposure unknown


37. Infectious syphilis case reports in BC by exposure category - total, 2006 to 2015



38. Infectious syphilis case reports in BC by exposure category - female, 2006 to 2015



39. Infectious syphilis case reports in BC by exposure category - male, 2006 to 2015



Infectious Syphilis among Men who have Sex with Men

Gay, bisexual, and other men who have sex with men (MSM) continue to be disproportionately affected by the current infectious syphilis epidemic in BC, constituting 84.4% (642/761 cases) of all cases in 2015 (Figure 37). The number of infectious syphilis cases among MSM has been steadily increasing since 2011. This is similar to the profile of syphilis epidemics in other Canadian provinces²⁰, the US²¹, and several European countries²² where syphilis cases were also predominantly among MSM and trends are increasing.

There were 642 cases among MSM in 2015, 21.3% (137 cases) were diagnosed with primary syphilis, 16.0% (103 cases) with secondary syphilis, and 62.6% (402 cases) were diagnosed with early latent infection. The proportion of infectious syphilis cases among MSM in the early latent stage of infection has steadily increased over time which may reflect increased syphilis testing. In 2015, 165 (25.7%) MSM cases had a prior syphilis diagnosis within the past five years, highlighting the importance of repeat infections in the current epidemic.²³

Despite recent increases in the number of annual cases, the characteristics of MSM with syphilis have remained similar over time. The number of cases increased for all age groups in 2015 (Figure 41) and the mean age of MSM diagnosed with syphilis was 40 years (range 17-82 years). In 2015, the majority of cases resided in the Lower Mainland with 71.7% (460 cases) residing in Vancouver and 16.4% (105 cases) in Fraser. As in previous years, the majority of cases in 2015 were among Caucasian (354 cases, 55.1%), Asian (52 cases, 8.1%), and Hispanic (39 cases, 6.1%) men (Table 40).


Living with HIV continues to be an important risk factor for acquiring infectious syphilis. In 2015, of the MSM cases with a known HIV status (620 cases), 51.3% (318 cases) were living with HIV at the time of their syphilis diagnosis which is a decrease from 2014 (265/447 cases, 59.3%) (Figure 42). The possible roles of core sexual networks and the biological synergy between HIV and syphilis are important areas of study which may help explain this trend. Coordinated public health follow-up, promotion of condom use, partner notification, and partner testing for all syphilis cases remain the cornerstone to controlling the syphilis epidemic for MSM in BC as do efforts to raise awareness among MSM. Given the continued increase of syphilis cases, the enhancement of ongoing programs and development of new syphilis control interventions for MSM remain a priority in BC.

40. Percentage of infectious syphilis case reports among MSM in BC by ethnicity, 2006 to 2015


Ethnicity	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
<i>No. Diagnoses</i>	157	174	229	146	115	154	310	476	466	642
Caucasian	79.6	73.0	75.5	68.5	75.7	70.1	64.8	59.9	55.4	55.1
Aboriginal	0.6	2.3	2.6	3.4	3.5	1.9	4.5	3.6	2.6	2.6
Asian	7.6	8.6	10.0	5.5	7.8	7.1	9.4	10.5	13.1	8.1
South Asian	3.2	1.7	3.1	0.0	3.5	5.8	3.2	2.7	2.4	2.2
Hispanic	6.4	8.0	4.8	13.0	3.5	9.7	7.1	5.7	7.1	6.1
Black	1.3	3.4	1.3	3.4	0.0	0.0	1.0	1.7	2.1	0.6
Other*	1.3	2.3	1.7	4.1	2.6	3.2	2.3	1.5	1.3	1.9
Unknown	0.0	0.6	0.9	2.1	3.5	1.9	7.7	14.5	16.1	23.4

* Other - Arab/West Asian and other/mixed ethnicity

41. Infectious syphilis case reports among MSM in BC by age group, 2006 to 2015



42. Infectious syphilis case reports* among MSM in BC by HIV co-infection, 2006 to 2015




Stage of Infection at Time of Syphilis Diagnosis

There have been slow changes in the proportion of infectious syphilis cases by stage of infection (i.e., primary, secondary, and early latent) over time. Most notably, the proportion of cases that are early latent syphilis has been increasing over time. This may reflect a greater uptake of syphilis testing or screening as people with syphilis infections in the early latent stage of infection are typically asymptomatic. In 2015, the proportion of cases that are early latent decreased slightly to 64.1% (488 cases) from 65.2% (358 cases) in 2014 (Figure 43). This increase may be a year-to-year variation.

From 2006 to 2015, 5.2% (195/3,759 cases) of infectious syphilis cases were also diagnosed with neurosyphilis. Neurosyphilis is commonly considered to be indicative of an advanced syphilis infection but can sometimes occur at an early stage of infection.

43. Stage of infection at time of syphilis diagnosis, 2006 to 2015




* Neurosyphilis is not mutually exclusive from other stages of syphilis infection

Maternal and Early Congenital Syphilis

Infectious syphilis acquired prior to or during pregnancy can be vertically transmitted to the infant which can have serious consequences, including infant death. Accordingly, prenatal screening for syphilis is recommended for all pregnant women in BC, as treatment will reduce the risk of transmission to or complications in infants.

In 2015, 51,131 syphilis tests were conducted as part of prenatal screening. Overall, the number of syphilis tests performed prenatally each year in BC is increasing slightly. There were no cases of congenital syphilis identified in BC in 2015 (Figure 44). In 2015, eleven maternal syphilis cases were reported whereas six cases were reported in 2014.

44. Maternal and early congenital syphilis case reports in BC, 2006 to 2015



* Rate per 100,000 live births

Endnotes

- ¹ For more information about the increase in chlamydia infections in high income countries around the world see: European Centre for Disease Prevention and Control. (February 2014). Chlamydia control in Europe: literature review. Retrieved from <http://www.ecdc.europa.eu/en/publications/Publications/chlamydia-control-europe.pdf>
- ² Reference to the increase in chlamydia infections in high income countries around the world see: Centers for Disease Control and Prevention, Division of STD Prevention. (October 2016). Sexually Transmitted Disease Surveillance 2015. Retrieved from <http://www.cdc.gov/std/stats15/std-surveillance-2015-print.pdf>
- ³ Reference to the “arrested immunity” hypothesis: Brunham RC, Rekart ML. (January 2008). The arrested immunity hypothesis and the epidemiology of chlamydia control. *Sexually Transmitted Diseases*, 35(1), 53-54.
- ⁴ Reference to the “arrested immunity” hypothesis: Rekart ML, Gilbert M, Meza R, Kim PH, Chang M, Money DM et al. (2012). Chlamydia public health programs and the epidemiology of pelvic inflammatory disease and ectopic pregnancy. *Journal of Infectious Diseases*, advance access published October 24, 2012. Retrieved from <http://jid.oxfordjournals.org/content/early/2012/10/24/infdis.jis644.full.pdf?keytype=ref&ijkey=y7YveY9ycdjbCr2>
- ⁵ For information about the increase in lymphogranuloma venereum (LGV) in the US: Schachter J, Moncada J. (June 2005). Lymphogranuloma venereum: how to turn an endemic disease into an outbreak of a new disease? Start looking. *Sexually Transmitted Diseases*, 32(6), 331-332. Retrieved from http://journals.lww.com/stdjournal/Citation/2005/06000/Lymphogranuloma_Venereum__How_to_Turn_an_Endemic.1.aspx
- ⁶ Reference to increase in lymphogranuloma venereum (LGV) in Europe: Childs T, Simms I, Alexander S, Eastick K, Hughes G, Field N. (December 3, 2015). Rapid increase in lymphogranuloma venereum in men who have sex with men, United Kingdom, 2003 to September 2015. *Eurosurveillance*, 20(48):pii=30076. Retrieved from <http://www.eurosurveillance.org/images/dynamic/EE/V20N48/art21320.pdf>
- ⁷ For more information about lymphogranuloma venereum (LGV) in BC: BC Centre for Disease Control. (March 2012). Lymphogranuloma venereum in British Columbia, 2004 to 2011. Retrieved from http://www.bccdc.ca/NR/rdonlyres/27E4F543-7D0E-417B-AF41-AEB262776FC3/0/STI_Reports_LGVinBC_20120404.pdf
- ⁸ For more information about lymphogranuloma venereum (LGV) in BC: BC Centre for Disease Control. (June 12, 2015). Lymphogranuloma venereum in BC: An update on a re-emerging STI. Retrieved from http://www.bccdc.ca/NR/rdonlyres/68FC8661-6622-42BE-8C71-F315382866E9/0/CPS_LGV_Update_20150612.pdf
- ⁹ For information about the how an individual with gonorrhea is at increased risk of acquiring HIV see: Wilton J. (Spring 2012). STIs: What role do they play in HIV transmission? CATIE, Canada’s source for HIV and hepatitis C information. Retrieved from <http://www.catie.ca/en/pif/spring-2012/stis-what-role-do-they-play-hiv-transmission>
- ¹⁰ Reference to how an individual with gonorrhea is at increased risk of acquiring HIV: Bernstein KT, Marcus JL, Nieri G, Philip SS, Klausner JD. (April 1, 2010). Rectal gonorrhea and chlamydia reinfection is associated with increased risk of HIV seroconversion. *Journal of Acquired Immune Deficiency Syndromes*, 53(4), 537-543. Retrieved from http://journals.lww.com/jaids/Fulltext/2010/04010/Rectal_Gonorrhea_and_Chlamydia_Reinfection_Is.16.aspx
- ¹¹ Reference to the increase in gonorrhea infections among MSM in other jurisdictions: Public Health England. (June 20, 2014). Sexually transmitted infections and chlamydia screening in England, 2013. Health Protection Report, 8(24). Retrieved from https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/326935/hpr2414.pdf
- ¹² Reference to the increase in gonorrhea infections among MSM in other jurisdictions: Wolitski RJ, Fenton KA. (2011). Sexual health, HIV, and sexually transmitted infections among gay, bisexual, and other men who have sex with men in the United States. *AIDS and Behavior*, 15 Suppl 1:S9-17.
- ¹³ For more information about the BC treatment guidelines for gonorrhea see: BC Centre for Disease Control. (revised August 2014). British Columbia Treatment Guidelines: Sexually Transmitted Infections in Adolescents and Adults 2014. Retrieved from http://www.bccdc.ca/resource-gallery/Documents/Communicable-Disease-Manual/Chapter%205%20-%20STI/CPS_BC_STI_Treatment_Guidelines_20112014.pdf

- ¹⁴ Reference to international surveillance data that suggests susceptibility of gonorrhea to cephalosporin treatments is threatened: Unemo M, Golparian D, Syversen G, Vestrheim DF, Moi H. (November 25, 2010). Two cases of verified clinical failures using internationally recommended first-line cefixime for gonorrhoea treatment, Norway, 2010. *Eurosurveillance*, 15(47). Retrieved from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19721>
- ¹⁵ Reference to international surveillance data that suggests susceptibility of gonorrhea to cephalosporin treatments is threatened: Yokoi S, Deguchi T, Ozawa T, Yasuda M, Ito S, Kubota Y, et al. (August 2007). Threat to cefixime treatment for gonorrhea. *Emerging Infectious Diseases*, 13(8), 1275-1277. Retrieved from: <http://wwwnc.cdc.gov/eid/content/13/8/pdfs/v13-n8.pdf>
- ¹⁶ Minimum inhibitory concentration (MIC) breakpoints to define “resistance” to cefixime and ceftriaxone have not yet been established, however, the Clinical and Laboratory Standards Institute (CLSI) defines MIC ≤ 0.25 $\mu\text{g/mL}$ as susceptible.
- ¹⁷ The US Centers for Disease Control and Prevention (CDC) has proposed MIC ≥ 2 $\mu\text{g/mL}$ as the non-susceptible threshold for azithromycin.
- ¹⁸ BC Stats. Census Statistical Profiles of Aboriginal Peoples, 2006. Retrieved from <http://www.bcstats.gov.bc.ca/statisticsbysubject/AboriginalPeoples/CensusProfiles.aspx>
- ¹⁹ See Endnote #18
- ²⁰ Reference to the increase in infectious syphilis cases among MSM in Canada: Public Health Agency of Canada. (2009). Executive Summary - Report on sexually transmitted infections in Canada: 2009. Retrieved from <http://www.phac-aspc.gc.ca/sti-its-surv-epi/sum-som-eng.php>
- ²¹ Reference to the increase in infectious syphilis cases among MSM in the US: See Endnote #2
- ²² Reference to the increase in infectious syphilis cases among MSM European countries: Savage EJ, Hughes G, Ison C, Lowndes CM, the European Surveillance of Sexually Transmitted Infections (ESSTI) network. (November 26, 2009). Syphilis and gonorrhea in men who have sex with men: a European overview. *Eurosurveillance*, 14(47). Retrieved from <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19417>
- ²³ For more information about the increase in infectious syphilis among MSM in BC: BC Centre for Disease Control. (June 2013). Infectious syphilis among gay, bisexual and other men who have sex with men in British Columbia, 2003 to 2012. Retrieved from http://www.bccdc.ca/NR/rdonlyres/B917A2F2-54C5-4691-8015-4F80538CAC1E/0/CPS_Report_Infectious_Syphilis_MSMBBC_20032012_20130624.pdf
- ²⁴ Clinical presentation includes any evidence of congenital syphilis on physical examination (e.g., hepatosplenomegaly), evidence of congenital syphilis on radiographs of long bones, a reactive CSF VDRL, an elevated CSF cell count or protein without other cause. Note that neonates may not display clinical manifestations of congenital syphilis and may meet laboratory criteria only.

Contributors

Epidemiology & Surveillance Team Clinical Prevention Services

Dr. Jason Wong, Physician Epidemiologist
Elsie Wong, Federal Field Surveillance Officer (PHAC)
Heming Jiang, Surveillance Analyst
Dr. Theodora Consolacion, Epidemiologist
Dr. Mark Gilbert, Medical Director

We would like to acknowledge the contributions of our many partners who without their support this report would not have been possible.

- Staff from the BCCDC Public Health Laboratory (PHL) for the collecting and compiling of gonorrhea antibiotic susceptibility data.
- Physicians, health care providers, and public health staff in BC for taking the time and effort to complete and submit case report forms.
- Staff from Clinical Prevention Services, BCCDC for the collecting (Epid and Syphilis Nurses) and entering (Clerical Team) of STI data.
- Aboriginal Health Physician Advisor from the Office of the Provincial Health Officer at the Ministry of Health and the First Nations Health Authority for providing feedback to the section pertaining to First Nations people.
- BC Ministry of Health for providing pelvic inflammatory disease and ectopic pregnancy data.
- BC Vital Statistics for providing live births data.
- Surveillance and Epidemiology Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada for providing the national chlamydia, gonorrhea and infectious syphilis rates.

Technical Appendix

Data Limitations

There are several key limitations to surveillance data which are important to understand in order to interpret surveillance data appropriately.

- Surveillance data are only reflective of the proportion of the population who test. Individuals with infections who have not tested are not included in the surveillance data presented in this report. Many people with sexually transmitted infections do not have symptoms and so do not seek health care advice or testing. This under-counting of cases may disproportionately affect some groups. For example, more women than men get syphilis testing in BC because of prenatal syphilis screening. Also, some sexually transmitted infections are more or less likely to show symptoms depending on sex and site of infection (e.g., urethral gonococcal infections in men are more likely to produce symptoms than those in women or those in the pharynx/throat).
- Surveillance trends are influenced by provider testing behaviours which may result in changes to the number of tests performed each year (e.g., increased vigilance for lymphogranuloma venereum in 2010-2011 may have resulted in more diagnoses of this disease, artificially driving up lymphogranuloma rates).
- Trends are influenced by temporal changes in testing technologies. Over the past ten years, nucleic acid amplification tests (NAAT) have replaced culture-based diagnostics for chlamydia and gonorrhea testing. The use of NAAT, which is a more sensitive test, has resulted in increased detection of these infections. NAAT technology has also allowed urine-based screening for chlamydia and gonorrhea thus reducing the use of urethral swabs for asymptomatic male clients. With the use of this less-invasive procedure, more men may be getting screened for chlamydia and gonorrhea.
- The majority of surveillance data presented in this surveillance report are extracted from case report forms completed by health care providers or public health nurses as part of the case follow-up process (which includes partner notification, patient education, and referral to appropriate services). There is an expected reporting delay to receipt of these forms which may range from days to months depending on the infection.
- Cases are classified by exposure category and ethnicity according to information elicited during follow-up from the case or their health care provider and, under-reporting of this information may lead to misclassification. For example, gay, bisexual and other men who have sex with men, and transgender persons may be reluctant to disclose these factors to their health care providers due to social stigma.
- Rates of infectious syphilis diagnoses among First Nations people are calculated with the numerator comprised of individuals with infectious syphilis who self identify as First Nations and the denominator comprised of individuals who are registered First Nations (see Data Sources below for further details).

Case Definitions

Sexually transmitted infections (STIs) included in this report are listed as reportable diseases in the *Communicable Disease Regulation* (Schedule A) of the *Public Health Act*.

Chlamydia

Genital: Detection and confirmation of *C. trachomatis* in anogenital (including rectal) or urinary specimens by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen).

Extra-genital: Detection and confirmation of *C. trachomatis* in specimens from the conjunctiva, pharynx and other extra-genital sites by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen).

Perinatally-acquired: Detection and confirmation of *C. trachomatis* in nasopharyngeal or other respiratory tract specimens by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen) from an infant who developed pneumonia in the first 6 months of life or conjunctival specimens from an infant who developed conjunctivitis in the first month of life.

Gonorrhea

Genital: Detection and confirmation of *N. gonorrhoeae* in anogenital (including rectal) or urinary specimens by appropriate laboratory techniques (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Extra-genital: Detection and confirmation of *N. gonorrhoeae* in specimens from the conjunctiva, pharynx, joint, blood, and other extra-genital sites by appropriate laboratory techniques (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Perinatally-acquired: Detection and confirmation of *N. gonorrhoeae* infection in the first 4 weeks of life leading to the diagnosis of gonococcal conjunctivitis, scalp abscess, vaginitis, bacteremia, arthritis,

meningitis, or endocarditis by an appropriate laboratory technique (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Infectious Syphilis

Infectious syphilis is a complex sexually transmitted infection that has a highly variable clinical course. Three stages of syphilis, primary, secondary and early latent, comprise infectious syphilis. Classification by a clinician with expertise in syphilis may take precedence over the following case definitions developed for surveillance purposes.

Primary Syphilis: Current clinical presentation compatible with primary syphilis (e.g., one or more ulcers/chancres), and one of the following:

- Identification of *T. pallidum* in clinical specimens (e.g., from chancre, regional lymph node) by dark field microscopy, direct fluorescent antibody, or nucleic acid amplification test (NAAT), or
- Reactive serology (treponemal, regardless of non-treponemal serology reactivity) in individuals with no previous history of syphilis, or
- Significant (i.e., four-fold or greater) increase in titre over the last known non-treponemal test.

Secondary Syphilis: Clinical presentation compatible with secondary syphilis (e.g., rash, fever, malaise, lymphadenopathy, mucus lesions, condyloma lata, alopecia, meningitis, headaches, uveitis, retinitis, recent hearing impairment), and one of the following:

- Identification of *T. pallidum* in clinical specimens (e.g., from chancre, regional lymph node) by dark-field microscopy, direct fluorescent antibody, or nucleic acid amplification test (NAAT), or

- Reactive serology (non-treponemal and treponemal) in individuals with no previous history of syphilis, or
- Significant (i.e., four-fold or greater) increase in titre over the last known non-treponemal test.

Early Latent Syphilis: An individual without symptoms of primary or secondary syphilis but has a reactive serology (non-treponemal and treponemal) or four-fold increase in titre over the last known non-treponemal test, and one of the following within the previous 12 months:

- Non-reactive serology, or
- Symptoms suggestive of primary or secondary syphilis, or
- Exposure to a sexual partner with primary, secondary or early latent syphilis, or
- Is a member of (or has had sexual partners in the previous 12 months from) groups at known increased risk of syphilis infection in BC, or
- Has a titre of $\geq 1:16$.

Early Congenital Syphilis: A stillbirth, neonate or older individual with clinical presentation²⁴ compatible with congenital syphilis, onset less than two years of age, and one of the following:

- Four-fold higher RPR than maternal titre and positive treponemal confirmatory test, or
- Detection of *T. pallidum* in clinical specimens (e.g., lesions, placenta, umbilical cord, autopsy) through darkfield microscopy, direct fluorescent antibody assay, or PCR, or
- Mother with untreated or inadequately treated syphilis (primary, secondary, early or late latent syphilis) during pregnancy or at birth.

Maternal Syphilis: A woman who meets the case definition of infectious syphilis (primary, secondary, early latent syphilis) or late latent syphilis, and one of the following:

- Syphilis serology conducted as part of prenatal blood screening, or
- Known to have given birth to an infant (live or stillborn) with congenital syphilis, or

- Clinical presentation with infectious syphilis during pregnancy.

Lymphogranuloma Venereum (LGV)

Confirmed: DNA sequencing for *C. trachomatis* confirming serovars of L1, L2, or L3 present.

Probable: One of the following two case definitions:

- Positive chlamydia NAAT or culture (from any site), and one of the following:
 - Proctitis, or
 - Inguinal/femoral lymphadenopathy, or
 - Suspicious lesion, or
 - Sexual partner who is confirmed or probable LGV case
- Clinical symptoms consistent with LGV (proctitis or inguinal/femoral lymphadenopathy or suspicious lesion) without a positive chlamydia test, and sexual partner who is confirmed or probable LGV case.

Data Sources

STI Data (Chlamydia, Gonorrhea, Infectious Syphilis)

When an individual is diagnosed with a reportable STI, the care provider completes a case report form (Health 208 form) then forwards it to BCCDC where the information is entered into the provincial STI database. Public health clinics with access to the provincial STI database directly enter the information for their newly diagnosed individuals.

Pelvic Inflammatory Disease and Ectopic Pregnancy Data

The diagnoses of pelvic inflammatory disease (PID) and ectopic pregnancy (EP) are captured in the Discharge Abstract Database (DAD) and the Medical Service Plan (MSP) payment database maintained by the BC Ministry of Health. The DAD includes data on patient discharges and day surgeries directly from hospitals in BC, including all known facilities for acute care and day surgery, and most facilities for

chronic care and rehabilitation. The MSP database contains data on insured medical services delivered on a fee-for-service basis. This data includes physician billings for inpatient and outpatient care, claims from supplementary health care practitioners, and claims for laboratory services and diagnostic procedures. MSP data do not include services provided via an Alternate Payment Program (e.g., contract or salary). Approximately 30% of BC physicians receive some remuneration through alternative payments. In particular, physicians in rural areas are more likely to be paid with Alternate Payment Programs. Some hospital emergency departments in BC are also funded through Alternate Payment Program contracts. Administrative data on PID and EP included in this surveillance report are extracted for women of reproductive age (15-44 years) who have at least one physician billing or hospital discharge per year based on the following International Classification of Disease (ICD) codes:

Pelvic Inflammatory Disease

ICD 9: MSP (1992-2010), DAD (1992-2000)

- Salpingitis and oophoritis (614, 614.0-614.2)
- Parametritis and pelvic cellulitis/peritonitis (614.3-614.5, 614.7)
- Other or unspecified inflammatory disease of female pelvic organs and tissues (614.8, 614.9)
- Inflammatory diseases of uterus except cervix (615, 615.0-615.9)

ICD 10: DAD (2001-2010)

- Salpingitis and oophoritis (N70, N70.0-N70.9)
- Parametritis and pelvic cellulitis/peritonitis (N73.0-N73.5)
- Other or unspecified female pelvic inflammatory diseases (N73.8, N73.9)
- Inflammatory diseases of uterus except cervix (N71, N71.0-N71.9)

Ectopic Pregnancy

ICD 9: MSP (1992-2010), DAD (1992-2000)

- Ectopic pregnancy (633)
- Abdominal pregnancy (633.0)
- Tubal pregnancy (633.1)

- Ovarian pregnancy (633.2)
- Other or unspecified ectopic pregnancy (633.8, 633.9)

ICD 10: DAD (2001-2010)

- Ectopic pregnancy (000)
- Abdominal pregnancy (000.0)
- Tubal pregnancy (000.1)
- Ovarian pregnancy (000.2)
- Other or unspecified ectopic pregnancy (000.8, 000.9)

BCCDC Public Health Laboratory (PHL)

Since July 2011, all rectal specimens that test positive for *C. trachomatis* in BC are routinely forwarded to the National Microbiology Laboratory in Winnipeg MB for LGV serovar testing via the BCCDC PHL. In addition, since 2012, clinics operated by the BCCDC have been routinely screening for *C. trachomatis* from extra-genital sites among those reporting behaviours that may put them at risk for pharyngeal or rectal infections.

The BCCDC PHL performs approximately 15-20% of all gonorrhea testing in the province, receiving specimens predominantly from Provincial Sexually Transmitted Infection Clinic sites operated by the BCCDC, from regional public health, youth, reproductive and sexual health clinics, and from hospitals throughout the province. At the BCCDC PHL, gonorrhea may be detected by nucleic acid amplification testing (NAAT) or conventional culture diagnostic methods. Culture testing is preferentially used for rectal and pharyngeal specimens and for all specimens from contacts to gonorrhea as well as patients who are symptomatic, not responding to treatment, or presenting for treatment after an initial NAAT-positive test. Antimicrobial susceptibility testing is routinely performed for all *N. gonorrhoeae* isolated by culture from clinical specimens. The BCCDC PHL additionally receives gonorrhea isolates forwarded for susceptibility testing from community or hospital-based laboratories in BC. Antimicrobial susceptibility testing is by E-test (bioMérieux) and data are analyzed by isolate.

Population Data

Unless noted otherwise, population data and associated rates are based on the P.E.O.P.L.E. 2015 Population Estimates and Projections released by BC Stats, BC Ministry of Technology, Innovation and Citizens' Services.

First Nations Population Estimates

Population rates for First Nations people are calculated using estimates from Aboriginal Affairs and Northern Development Canada.

These estimates are based on the Indian Registry System (IRS), which includes individuals who have registered for First Nations status under the Indian Act. The IRS is subject to several limitations, including:

- Under-counting due to delayed reporting of infants entitled to be registered, as well as other unregistered individuals who are entitled for status designation
- Over-counting due to individuals remaining on the IRS after they are deceased
- Geographic misclassification because individuals are included in the BC population according to membership of a BC band rather than current place of residence
- Systematic biases from imbalance in the migration into and out of the BC region (these are difficult to quantify)

For further details about the data source and its limitations, see the report entitled *Registered Indian Population by Sex and Residence, 2014: Statistics and Measurement Directorate (2014)*. Aboriginal Affairs and Northern Development Canada.

Live Births

Perinatal rates are calculated using live births data from the BC Vital Statistics Agency.

Additional Notes

Classification of Health Region

Cases are assigned to health regions (i.e., Health Authority or Health Service Delivery

Area) by residence. If residence is unknown, the case is then assigned to the health region where the individual was tested.

Classification of Ethnicity

Ethnicity is based on information elicited from the case or health care provider during follow-up. Since ethnicity data for chlamydia and gonorrhea cases are often not collected they are not included in this report.

Ethnicity	Example
Aboriginal	First Nations, Inuit, Métis
Arab/West Asian	Armenian, Egyptian, Iranian, Moroccan, Lebanese, Afghani
Asian	Chinese, Japanese, Vietnamese, Cambodian, Indonesian, Filipino, Korean, Laotian
Black	African, Haitian, Jamaican, Somali
Caucasian (White)	Irish, Scottish, English, Portuguese, Italian, Russian
Hispanic	Mexican, Central/South American
South Asian	East Indian, Pakistani, Sri Lankan, Punjabi, Bangladeshi
other/mixed ethnicity	ethnicity is known but is not included in one of the above categories or case has dual ethnicity
unspecified	information about ethnicity is not elicited from case or health care provider

Exposure Group Hierarchy

Cases may have more than one type of sexual exposure. The following are definitions of sexual exposures used in this surveillance report. For infectious syphilis cases, individuals are assigned to the exposure category listed first (or highest) in the following hierarchy.

1. **MSM***: Male who reports having male sex partner(s), with or without female sex partners.
2. **Street-Involved, Sex Trade Worker and Patron**:
 - i) Street-Involved – Person who reports either: (a) living on the street or in a single room occupancy (SRO) hotel;

- or (b) attached to the street; or (c) having no fixed address; or (d) transient
 - ii) Sex Trade Worker (STW) – Person who reports providing sex to another individual in exchange for money, shelter, food, drugs, etc.
 - iii) Patron of STW – Person who reports payment (with money, shelter, food, drugs, etc.) for sex with a STW
3. **Heterosexual Contact***: Male who reports having female sex partner(s) only or female who reports having male with/without female sex partner(s).
4. **Acquired Outside of Canada:**
- i) Foreign Acquired – Person currently residing in Canada but likely acquired syphilis outside of Canada (i.e., reports sexual partner(s) in other countries)
 - ii) Immigration – Person immigrating to Canada and identified with syphilis through testing done as part of the immigration process
5. **Other Risk Factor:** Likely route of exposure is known but cannot be classified into any of the major exposure categories listed here. For example, females reporting female sex partner(s) only.
6. **Unknown:** Route of exposure is unknown or not identified at the time of completion of case follow-up (e.g., route of exposure not provided by case).

* A transgender individual may be assigned to either MSM or Heterosexual Contact exposure category depending on how this individual describes their sexual partners.