2011-12: Number 16, Week 16 April 15 to 21, 2012



Prepared by BCCDC Influenza & Emerging Respiratory Pathogens Team

Continued low-level influenza activity in BC

		Contents:		
British Columbia:			International:	Page 7
Sentinel Physicians	Page 2			
Children's Hospital ER	Page 2		Other:	
Medical Services Plan	Page 3		List of Acronyms	Page 8
Laboratory Surveillance	Page 5		Web Sites	Page 8
ILI Outbreaks	Page 6		Outbreak Report Form	Page 9
Canada:				
FluWatch Activity levels	Page 6			
NML Strain Characterization	Page 6			
NML Antiviral Resistance	Page 7			

Summary

In week 16 (April 15-21, 2012), most influenza surveillance indicators suggested that influenza activity continues at low levels in BC. The proportion of patients with influenza-like illness among those presenting to sentinel physicians was 0.20%, similar to previous weeks and within the expected range for this time of year. Throughout the province, influenza illness as a proportion of all submitted BC MSP claims remained at or below the 10-year median for this time of year. In week 16, no lab-confirmed influenza outbreaks were reported; one has been reported in week 17 (influenza A, subtype pending). Of the one hundred and six specimens tested at the BC Public Health Microbiology & Reference Laboratory, PHSA, during this period, 18 (17.0%) were positive for influenza, including 8 (7.5%) influenza A/H3N2, 3 (2.8%) A(H1N1)pdm09, 2 (1.9%) influenza A (subtype pending), and 5 (4.7%) influenza B. Other significant respiratory virus detections included rhino/enterovirus (13/106, 12.3%), respiratory syncytial virus (12/106, 11.3%), and human bocavirus (8/106, 7.5%). Other respiratory viruses were also sporadically detected. RSV increased and continued to dominate among the respiratory viruses detected at BC Children's Hospital.

Report disseminated April 26, 2012 Contributors: Lisan Kwindt, Naveed Janjua, Danuta Skowronski


2011-12: Number 16, Week 16 April 15 to 21, 2012

British Columbia

Sentinel Physicians

In week 16, the proportion of patients with ILI among those presenting to sentinel physicians was 0.20%, similar to the previous week and within the expected range for this time of year. To date, 59% of sentinel physician sites have reported for week 16.

Percentage of Patient Visits due to Influenza Like Illness (ILI) per Week Compared to Average Percentage of ILI Visits for the Past 20 Seasons Sentinel Physicians, British Columbia, 2011-2012



^{*} Data subject to change as reporting becomes increasingly complete.

BC Children's Hospital Emergency Room

The percentage of BC Children's Hospital ER visits attributed to "fever and cough" or flu-like illness is unavailable pending upgrades to the data collection system.


[†]Historical values exclude 2008-09 and 2009-10 seasons due to atypical seasonality.

2011-12: Number 16, Week 16 April 15 to 21, 2012

Medical Services Plan

In week 16, influenza illness as a proportion of all submitted BC Medical Services Plan (MSP) claims remained low, at or below the 10-year median level for this time of year throughout the province.


Influenza Illness Claims* British Columbia



^{*} Influenza illness is tracked as the percentage of all submitted MSP general practitioner claims with ICD-9 code 487 (influenza). Data provided by Population Health Surveillance and Epidemiology, BC Ministry of Health Services

Note: MSP week beginning 28 August 2011 corresponds to sentinel ILI week 35; Data current to 25 April 2012

Northern




2011-12: Number 16, Week 16 April 15 to 21, 2012


Interior

3.0% Above historical maximum Above 10-year maximum Above 10-year 75th percentile Between 25th & 75th percentile 2.5% Below 10-year 25th percentile Below 10-year minimum 2.0% 10-year median 2009/2010 H1N1 pandemic yea 1.5% 1.0% 0.5% 0.0% 15 Aug, 2011 29 Aug, 2011 12 Sep, 2011 26 Sep, 2011 10 Oct, 2011 Oct, 2011 Nov, 2011 19 Dec, 2011 2012 2012 2 Jan, 2012 16 Jan, 2012 30 Jan, 2012 13 Feb, 2012 27 Feb, 2012 12 Mar, 2012 26 Mar, 2012 7 May, 2012 1 May, 2012 4 Jun, 2012 18 Jun, 2012 21 Nov, 2011 5 Dec, 2011 9 Apr, 2012 23 Apr, 2012


Vancouver Coastal



Fraser




Vancouver Island




2011-12: Number 16, Week 16 April 15 to 21, 2012

Laboratory Reports


In week 16, one hundred six specimens were tested for influenza viruses at the BC Public Health Microbiology & Reference Laboratory, PHSA. Among them, eighteen (17.0%) were positive for influenza viruses, including 8 (7.5%) influenza A/H3N2 from VCHA, VIHA, and NHA; 3 (2.8%) A(H1N1)pdm09 from FHA, VCHA, and VIHA; 2 (1.9%) influenza A (subtype pending); and 5 (4.7%) influenza B from VIHA. Of 106 specimens tested for other respiratory viruses, significant detections included rhino/enterovirus (13/106, 12.3%), respiratory syncytial virus (12/106, 11.3%), and human bocavirus (8/106, 7.5%). Other respiratory viruses were also sporadically detected.





In week 16, BC Children's and Women's Health Centre Laboratory tested 51 respiratory specimens: 4 (7.8%) were positive for influenza virus, slightly higher than the preceding week, including 2 influenza A and 2 influenza B. RSV increased and continued to predominate among respiratory viruses detected (15/51, 29.4%). Other respiratory viruses were also detected at low levels.

Influenza and Other Virus Detections Among Respiratory Specimens Submitted to BC Children's and Women's Health Centre Laboratory, 2011-2012




Data provided by Virology Department at Children's & Women's Health Centre of BC

2011-12: Number 16, Week 16 April 15 to 21, 2012

ILI Outbreaks

In week 16, no lab-confirmed influenza outbreaks were reported. One lab-confirmed influenza A outbreak was reported in week 17 in VIHA (subtype pending).

> Number of Influenza and Influenza-Like Illness (ILI) Outbreaks Reported, Compared to Current Sentinel ILI Rate and Average Sentinel ILI Rate for past 20 years, per Week, British Columbia, 2011-2012 season



^{*} Facility influenza outbreak defined as 2 or more ILI cases within 7-day period, with at least one case laboratory-confirmed as influenza.

CANADA

FluWatch

In week 15 (April 8 to April 14, 2012), influenza activity in Canada continued to decline; however, activity remained elevated in some regions of the country (i.e. Atlantic Region, Ontario, Alberta & British Columbia). In total 841 laboratory detections of influenza were reported in week 15: 13.9% A/H3N2, 6.2% A(H1N1)pdm09, 16.5% un-subtyped influenza A and 63.4% influenza B. The ILI consultation rate in week 15 declined slightly compared to the previous week and remained within the expected levels for this time of year. PHAC further reported 118 laboratory-confirmed influenza-associated hospitalizations including 24 paediatric (two thirds due to influenza B, and one third due to influenza A) and 94 adult (69% due to influenza B, and 31% due to influenza A). www.phac-aspc.gc.ca/fluwatch/

National Microbiology Laboratory (NML): Strain Characterization

Between September 1, 2011 and April 26, 2012, 969 isolates were collected from provincial and hospital labs and characterized at the NML as follows:

- 189 A/Perth/16/2009-like (H3N2)1 from NS, QUE, ONT, MAN, SASK, ALTA, BC, and NT;
- 178 A/California/07/09-like (H1N1)* from NB, QUE, ONT, MAN, SASK, ALTA, and BC;
- 306 B/Brisbane/60/2008-like (B/Victoria/02/87 lineage)[†] from NFLD, NS, NB, QUE, ONT, MAN, SASK, ALTA, and BC:

296 B/Wisconsin/01/2010-like (recent B Yamagata lineage) from NS, NB, QUE, ONT, MAN, SASK, ALTA, BC, and NU:

[†] School ILI outbreak defined as >10% absenteeism on any day, most likely due to ILI.
** Historical values exclude 2008-09 and 2009-10 seasons due to atypical seasonality.

¹ indicates a strain match to the recommended H3N2 component of the 2011-12 northern hemisphere influenza vaccine

^{*} indicates a strain match to the recommended H1N1 component for the 2011-2012 northern hemisphere influenza vaccine

[†] indicates a strain match to the recommended influenza B component for the 2011-2012 influenza vaccine

2011-12: Number 16, Week 16 April 15 to 21, 2012

NML: Antiviral Resistance

From September 1, 2011 to April 26, 2012, drug susceptibility testing was performed at the NML for influenza A/H3N2 (oseltamivir: 182; zanamivir: 183; amantadine: 309), influenza A(H1N1)pdm09 (oseltamivir: 190; zanamivir: 190; amantadine: 249), and influenza B isolates (oseltamivir: 565; zanamivir: 565). The results indicated that all isolates were sensitive to oseltamivir and zanamivir, while all influenza A/H3N2 isolates but one, and all A(H1N1)pdm09 isolates, were resistant to amantadine.

INTERNATIONAL

<u>USA:</u> In week 15, ending 14 April 2012, influenza activity in the United States was elevated in some areas, but declined nationally and in most regions. Six hundred and fifty-three (17.5%) specimens tested were positive for influenza, including 533 (81.6%) influenza A [289 A/H3N2, 106 A(H1N1)pdm09, and 138 un-subtyped A] and 120 (18.4%) influenza B. The proportion of outpatient visits for ILI was 1.5% which was below the national baseline of 2.4%. The proportion of all deaths due to pneumonia and influenza illness was 7.0%, below the epidemic threshold of 7.7% for week 15. Two influenza-associated paediatric deaths were reported to the US CDC during week 15, one associated with A(H1N1)pdm09, and the other with A(H3N2). www.cdc.gov/flu/weekly/.

WHO news: (last updated on 13 April 2012). In most of the northern hemisphere temperate regions, influenza activity had peaked and was declining. In North America, influenza indicators remained elevated in some areas of the United States of America, but declined in the last couple of weeks. In Europe and northern Asia, nearly every country had passed its peak of transmission and reported declining activity. The most commonly detected virus type or subtype throughout Europe and North America (except Mexico) had been influenza A(H3N2), although the proportion of influenza B detection had been increasing toward the end of the season in North America. In Mexico influenza A(H1N1)pdm09 had been the most common influenza virus circulating; China and the surrounding countries of northern Asia were still reporting a predominance of influenza type B virus. No significant change in antiviral resistance was reported so far this season.

www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html

Avian Influenza:

No new cases of avian influenza have been reported by the WHO since the last update (12 April 2012). www.who.int/influenza/human_animal_interface/avian_influenza/en/

WHO Recommendations for 2012-13 Northern Hemisphere Influenza Vaccine

On 23 February, 2012 the WHO announced the recommended strain components for the 2012-13 northern hemisphere vaccine:

A/California/7/2009 (H1N1)pdm09 virus A/Victoria/361/2011 (H3N2)-like virus*

B/Wisconsin/1/2010 (Yamagata lineage)-like virus*

* these two of the three recommended components are different from the northern hemisphere seasonal TIV vaccines produced and administered in 2010-11 and 2011-2012. For further details, see: www.who.int/influenza/vaccines/virus/recommendations/2012 13 north/en/index.html

2011-12: Number 16, Week 16 April 15 to 21, 2012

Contact Us:

Communicable Disease Prevention and Control (CDPACS):
BC Centre for Disease Control (BCCDC)

List of Acronyms

ACF: Acute Care Facility
AI: Avian influenza

FHA: Fraser Health Authority **HBoV**: Human bocavirus **HMPV**: Human metapneumovirus

HSDA: Health Service Delivery Area
IHA: Interior Health Authority
ILI: Influenza-Like Illness
LTCF: Long-Term Care Facility

VIF WF

VCHA: Vancouver Coastal Health Authority VIHA: Vancouver Island Health Authority WHO: World Health Organization

NML: National Microbiological Laboratory

MSP: BC Medical Services Plan

NHA: Northern Health Authority

pH1N1: Pandemic H1N1 influenza

RSV: Respiratory syncytial virus

Web Sites

1. Influenza Web Sites

Canada – Flu Watch: www.phac-aspc.gc.ca/fluwatch/
Washington State Flu Updates: www.doh.wa.gov/FLUNews/
USA Weekly Surveillance reports: www.cdc.gov/flu/weekly/
European Influenza Surveillance Scheme: www.ecdc.europa.eu

WHO - Global Influenza Programme: www.who.int/csr/disease/influenza/mission/

WHO – Weekly Epidemiological Record: www.who.int/wer/en/

Influenza Centre (Australia): www.influenzacentre.org/

Australian Influenza Report: www.health.gov.au/internet/main/publishing.nsf/content/cda-surveil-ozflu-flucurr.htm
New Zealand Influenza Surveillance Reports: www.surv.esr.cri.nz/virology/influenza weekly <a href="www.surv.esr.cri.nz/virology/influenza weekly <a href="www.surv.esr.cri.nz/virology/influenza weekly <a href="www.surv.esr.cri.nz/virology/influenza weekly <a href="www.surv.esr.cri.nz/virology/

2. Avian Influenza Web Sites

World Health Organization – Avian Influenza: www.who.int/csr/disease/avian influenza/en/ World Organization for Animal Health: www.oie.int/eng/en_index.htm

3. This Report On-line: www.bccdc.ca/dis-cond/DiseaseStatsReports/influSurveillanceReports.htm

version: 26 Oct 2011

Phone: (604) 707-2510

Fax: (604) 707-2516

ilioutbreak@bccdc.ca

Influenza-Like Illness (ILI) Outbreak Summary Report Form

Please complete and email to ilioutbreak@bccdc.ca

Note: This form is for provincial surveillance purposes.

Please notify your local health unit per local guidelines/requirements.

ILI: Acute onset of respiratory illness with fever and cough and with one or more of the following: sore throat, arthralgia, myalgia, or prostration which could be due to influenza virus. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent. Schools and work site outbreak: greater than 10% absenteeism on any day, most likely due to ILI. Residential institutions (facilities) outbreak: two or more cases of ILI within a seven-day period. Health unit/medical health officer notified? ☐ Yes ☐ No Reporting Information Person Reporting: _____ Title: _____ Contact Phone: _____ Email: _____ Health Authority: _____ HSDA:_____ Full Facility Name: First Notification (complete section **B** below: Section **D** if available) Is this report: Update (complete section **C** below; Section **D** if available) Outbreak Over (complete section **C** below; Section **D** if available) **First Notification** B Type of facility: Senior's Residence Acute Care Hospital (if ward or wing, please specify name/number: Date of onset of first case of ILI (dd/mm/yyyy): __DD__/_MMM__/_YYYY Residents/Students Numbers to date Staff Total With ILI Hospitalized Died **Update AND Outbreak Declared Over** Date of onset for most recent case of ILI (dd/mm/yyyy): __DD_/_MMM_/_YYYY If over, date outbreak declared over (dd/mm/yyyy): DD / MMM / YYYY Numbers to date Residents/Students Staff Total With ILI Hospitalized Died **Laboratory Information** Specimen(s) submitted? ☐ Yes (location: _____) ☐ No ☐ Don't know