

**COVID-19 mRNA Vaccine  
COMIRNATY®  
OMICRON LP.8.1  
(5-11 years of age presentation: 10 mcg/0.3 mL)**

**Supplier: Pfizer**  
**Light Blue Vial Cap**  
**Light Blue Label Border**

**INDICATIONS:**

- Individuals 5-11 years of age (inclusive). See [COVID-19 Vaccine Eligibility](#).

The vaccine is approved for use in those 5-11 years of age only.

**DOSES AND SCHEDULE:<sup>A</sup>**

- Individuals 5-11 years of age: 1 dose given as 0.3 mL (10 mcg) **IM** at least 3 months <sup>B</sup> after last dose of COVID-19 vaccine, regardless of previous COVID-19 vaccination history.
- Individuals 5-11 years of age who are moderately to severely immunosuppressed:  
Individuals in this age group who are moderately to severely immunosuppressed (see [COVID-19 Vaccine Eligibility](#)) should receive a total of at least 2 doses of COVID-19 vaccine per age-appropriate dosing recommendation above, with at least one of these doses being the COVID-19 LP.8.1 formulation. Refer to intervals in table below. Those who self-declare a recommendation by their health care provider are eligible to receive a 3-dose series, at a recommended interval of 8 weeks between doses.<sup>C</sup> New recipients of HSCT or CART should receive a 3-dose series, at a recommended interval of 8 weeks between doses.

Recommendations for Moderately to Severely Immunosuppressed Clients		
COVID-19 Vaccination History	Number of Dose(s) of COVID-19 LP.8.1 Vaccine	Recommended Interval Between Doses
2 or more doses	1 dose	3 months after last dose <sup>B</sup>
1 dose	1 dose	8 weeks after last dose <sup>D</sup>
0 doses	2 doses	8 weeks between doses <sup>D</sup>

**ADMINISTRATION:**

- The vaccine is supplied as a 0.3 mL single dose vial (SDV) that does not require dilution or reconstitution.

**Storage and Handling:<sup>E</sup>**

**Storage:**

- Single dose vials of vaccine may arrive frozen at ultra-low temperatures (-90°C to -60°C) in thermal containers with dry ice or at refrigerated temperatures (+2°C to +8°C).

<sup>A</sup> Children who initiated COVID-19 vaccination at 4 years of age and turn 5 years of age prior to completion of primary series should complete their primary series based on the age of series initiation. See Part 4 - Biological Products, [COVID-19 Vaccine Eligibility](#).

<sup>B</sup> An interval of at least 3 months can be used due to the unknown seasonality of SARS-CoV-2 and the expected duration of protection from vaccine.

<sup>C</sup> Additional doses above the authorized schedule are intended to improve the immune response in individuals who are moderately to severely immunosuppressed. This may be especially important in individuals who have not had a prior SARS-CoV-2 infection as they do not benefit from hybrid immunity. For such individuals, a total of 3 doses may be provided to enhance protection.

<sup>D</sup> A 28 day minimum interval may be considered. For optimal response, the recommended interval should be observed.

<sup>E</sup> For more information on storage and handling and temperature monitoring refer to [Appendix E: Management of Biologicals](#) and [Guidance for Receiving and Handling the Pfizer-BioNTech COVID-19 mRNA Vaccine \(including dry ice procedures\)](#).

## COVID-19 mRNA Vaccine

COMIRNATY®

OMICRON LP.8.1

(5-11 years of age presentation: 10 mcg/0.3 mL)

**Supplier: Pfizer**

Light Blue Vial Cap

Light Blue Label Border

### ADMINISTRATION (continued):

- Frozen vaccine:
  - Vaccine that arrives frozen can be stored in an ultra-low temperature freezer at -90°C to -60°C up to the end of its expiry date and kept in the original packaging, until ready to use.
  - **Do not store vaccine at -25°C to -15°C.**
  - The vaccine can be stored for up to 30 days in a validated thermal container with dry ice; requires re-icing with 20-23 kg of new dry ice every 5 days if opened twice daily.
- Refrigerated vaccine:
  - Vaccine that arrives under refrigerated conditions should be stored refrigerated (+2°C to +8°C). Do not freeze vaccine.
  - Vaccine may be stored refrigerated for up to 10 weeks from the date of transfer from frozen storage to refrigerated storage conditions. Ensure the vaccine carton has been updated to reflect the 10-week expiry date.

### Thawing:

- Single dose vials of frozen vaccine can be thawed in a refrigerator (+2°C to +8°C) or at room temperature (up to +25°C).
  - From the freezer to the refrigerator:
    - A carton of 10 vials may take up to 2 hours to thaw in a refrigerator (+2°C to +8°C).
    - Once moved to a refrigerator, the vaccine may be stored at +2°C to +8°C for up to 10 weeks. Ensure that the 10-week expiry date is included on the vaccine carton.
  - From the freezer to room temperature:
    - Vials can be thawed at room temperature (up to +25°C) for 30 minutes. The vaccine can be stored at room temperature (up to +25°C) for up to 12 hours prior to use.
  - **Do not refreeze thawed vials.**

### Use:

- Prior to use and after thawing, gently invert the vial 10 times to mix; **do not shake**.
- Prior to mixing, the thawed vaccine is a clear to slightly opalescent suspension and may contain white to off-white opaque amorphous particles.
- After mixing the vaccine should appear as a clear to slightly opalescent suspension with no visible particles. Do not use if liquid is discolored or if particles are observed after mixing.

---

### BOOSTER DOSES:

No further doses are recommended following COVID-19 vaccination with the LP.8.1 formulation at this time.

**COVID-19 mRNA Vaccine  
COMIRNATY®  
OMICRON LP.8.1  
(5-11 years of age presentation: 10 mcg/0.3 mL)**

**Supplier: Pfizer**  
**Light Blue Vial Cap**  
**Light Blue Label Border**

**SEROLOGICAL TESTING:**

Serological testing is not recommended before or after immunization.

---

**CONTRAINDICATIONS:**

1. History of anaphylactic reaction to any component of the vaccine is generally considered a contraindication, however for more details on the administration of COVID-19 vaccines to individuals with allergies to components of the COVID-19 vaccines, please see the PRECAUTIONS section.

---

**PRODUCT COMPONENTS:**

Potential allergens: 2-[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide.

Other components: ((4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate); 1,2-distearoyl-sn-glycero-3-phosphocholine; cholesterol; sucrose; tromethamine, tromethamine hydrochloride.

---

**PRECAUTIONS:**

- For individuals with a history of anaphylactic reaction to a previous dose of an mRNA COVID-19 vaccine, re-vaccination (i.e., administration of a subsequent dose in the series when indicated) may be offered with an mRNA vaccine if a risk assessment deems that the benefits outweigh the potential risks for the individual and if informed consent is provided. Prior to re-vaccination, consultation with an allergist or another appropriate physician (e.g., Medical Health Officer) is advised. If re-vaccinated, vaccine administration should be done in a controlled setting with expertise and equipment to manage anaphylaxis, with an extended period of observation of at least 30 minutes after re-vaccination.
- For individuals with suspected hypersensitivity or non-anaphylactic allergy to COVID-19 vaccine components, the vaccine should be administered in a controlled setting with expertise and equipment to manage anaphylaxis, with an extended period of observation post-vaccination of at least 30 minutes.
- Wait until symptoms of an acute illness are resolved before vaccinating with COVID-19 vaccine to differentiate symptoms of illness from vaccine side effects.
- Individuals diagnosed with Multisystem Inflammatory Syndrome in Children (MIS-C) should delay COVID-19 vaccination until they have recovered from illness and for 90 days after the date of diagnosis of MIS-C.
- Additional doses of a COVID-19 vaccine should be deferred in individuals who experienced a physician-diagnosed myocarditis or pericarditis event following a previous dose of a COVID-19 vaccine. However, those with a history compatible with pericarditis who had no cardiac workup or had normal investigations may proceed to further vaccination once symptoms have resolved and 90 days have passed since receipt of the dose associated with the event. For those with confirmed myocarditis (with or without pericarditis), an individual risk/benefit discussion between the patient and their healthcare provider should occur so that the patient (with their parent/guardian as applicable) can make an informed decision about proceeding with a subsequent dose. If another dose is offered, the Pfizer-BioNTech COVID-19 vaccine should be offered because of the lower rate of myocarditis/pericarditis compared to the Moderna vaccine. Informed consent should include the unknown rates of recurrence of myocarditis and/or pericarditis following receipt of

## COVID-19 mRNA Vaccine

**COMIRNATY®**

**OMICRON LP.8.1**

**(5-11 years of age presentation: 10 mcg/0.3 mL)**

**Supplier: Pfizer**

**Light Blue Vial Cap**

**Light Blue Label Border**

### **PRECAUTIONS (continued):**

additional doses of the Pfizer-BioNTech COVID-19 vaccine, as well as the need to seek immediate medical assessment and care should symptoms develop. Deferral is not required for those with a prior history of myocarditis or pericarditis that is unrelated to COVID-19 vaccines and are no longer being followed by a medical professional for heart issues. For more information refer to the [NACI](#) summary.

---

### **SPECIAL CONSIDERATIONS:**

- COVID-19 vaccines can be administered concomitantly or at any time before or after the administration of another inactivated or live vaccine.
- Deferral of COVID-19 vaccination is no longer recommended for individuals who have received anti-SARS-CoV-2 monoclonal antibodies or convalescent plasma for treatment or prevention of COVID-19.
- COVID-19 vaccine may be offered to individuals without contraindications who have recovered from confirmed SARS-CoV-2 infection.
- For previously vaccinated individuals, COVID-19 vaccine may be deferred in those who have tested positive for COVID-19 (by PCR or rapid antigen test) until 3 months from symptom onset or, for asymptomatic cases, from the time of the positive test.
- For individuals that have not yet completed their primary series, COVID-19 vaccine may be deferred in those who have tested positive for COVID-19 (by PCR or rapid antigen test) until 8 weeks from symptom onset or, for asymptomatic cases, from the time of the positive test. If these individuals are moderately to severely immunosuppressed, a 4-8 week interval may be considered.

---

### **ADVERSE EVENTS:**

**Local:** pain, swelling, redness.

**Systemic:** fatigue, headache, myalgia, chills, arthralgia, fever, nausea and vomiting.

Rare cases of facial paralysis/Bell's palsy have been reported.

Rare cases of myocarditis (inflammation of the heart muscle) and/or pericarditis (inflammation of the lining around the heart) have been reported following vaccination with prior formulations of COVID-19 vaccines. These are more often seen after the 2<sup>nd</sup> dose received, when spacing between the first and second dose was less than 8 weeks, in males, in those 12-29 years old, and with use of a Moderna 100 mcg dose (used in primary series but not in subsequent booster doses). Typical onset of symptoms is within the first week after vaccination. This event has not been observed in younger children. Follow-up studies of those who experienced myocarditis continues, but the majority of individuals respond well to conservative therapy such as anti-inflammatory treatment and recover quickly.